

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
29 July 2004 (29.07.2004)

PCT

(10) International Publication Number  
WO 2004/062665 A1

(51) International Patent Classification<sup>7</sup>: A61K 31/437,  
A61P 1/00, 25/18, 25/22, 25/24, 29/00, C07D 471/04

Alessandro Fleming 2, I-37100 Verona (IT). ST-DENIS,  
Yves [CA/IT]; GlaxoSmithKline SpA, Via Alessandro  
Fleming 2, I-37100 Verona (IT).

(21) International Application Number:

PCT/EP2004/000409

(74) Agent: GIDDINGS, Peter, John; GlaxoSmithKline,  
CN925.1, Corporate Intellectual Property, 980 Great West  
Road, Brentford Middlesex TW8 9GS (GB).

(22) International Filing Date: 14 January 2004 (14.01.2004)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

60/440,432

16 January 2003 (16.01.2003) US

(71) Applicants (for all designated States except US): SB  
PHARMCO PUERTO RICO INC [US/US]; The United  
States Corporation Company, 105 Ponce de Leon Avenue,  
One Comptroller Plaza, Hato Rey, Puerto Rico 00917  
(US). NEUROCRINE BIOSCIENCES INC [US/US];  
10555 Science Center Drive, San Diego, CA 92121 (US).

(81) Designated States (unless otherwise indicated, for every  
kind of national protection available): AB, AG, AL, AM,  
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,  
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,  
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,  
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,  
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,  
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,  
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,  
ZW.

(71) Applicant (for MG only): GLAXO GROUP LIMITED  
[GB/GB]; Glaxo Wellcome House, Berkeley Avenue,  
Greenford, Middlesex UB6 0NN (GB).

(84) Designated States (unless otherwise indicated, for every  
kind of regional protection available): ARIPO (BW, GH,  
GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), Euro-  
pean (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR,  
GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK,  
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,  
ML, MR, NE, SN, TD, TG).

(72) Inventors; and


(75) Inventors/Applicants (for US only): CASTIGLIONI,  
Emiliano [IT/IT]; GlaxoSmithKline SpA, Via Alessandro  
Fleming 2, I-37100 Verona (IT). DI FABIO, Romano  
[IT/IT]; GlaxoSmithKline SpA, Via Alessandro Fleming  
2, I-37100 Verona (IT). FERLANI, Aldo [IT/IT]; Glaxo-  
SmithKline SpA, Via Alessandro Fleming 2, I-37100  
Verona (IT). MICHELI, Fabrizio [IT/IT]; GlaxoSmithK-  
line SpA, Via Alessandro Fleming 2, I-37100 Verona (IT).  
SABBATINI, Fabio [IT/IT]; GlaxoSmithKline SpA, Via

Published:

- with international search report
- before the expiration of the time limit for amending the  
claims and to be republished in the event of receipt of  
amendments

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: HETEROARYL- SUBSTITUTED PYRROLO' 2, 3- B' PYRIDINE DERIVATIVES AS CRF RECEPTOR ANTAGO-  
NISTS



(57) Abstract: The present invention provides compounds of formula (I) including stereoisomers, prodrugs and pharmaceutically acceptable salts or solvates thereof, to processes for their preparation, to pharmaceutical compositions containing them and to their use in the treatment of conditions mediated by corticotropin-releasing factor (CRF).

WO 2004/062665 A1

## Chemical compounds

The present invention relates to bicyclic derivatives, to processes for their preparation, to pharmaceutical compositions containing them and to their use in therapy.

The first corticotropin-releasing factor (CRF) was isolated from ovine hypothalamus and identified as a 41-amino acid peptide (Vale et al., Science 213: 1394-1397, 1981).

CRF has been found to produce profound alterations in endocrine, nervous and immune system function. CRF is believed to be the major physiological regulator of the basal and stress-release of adrenocorticotrophic hormone ("ACTH"), Bendorphin and other proopiomelanocortin ("POMC")-derived peptides from the anterior pituitary (Vale et al., Science 213: 1394-1397, 1981).


In addition to its role in stimulating the production of ACTH and POMC, CRF appears to be one of the pivotal central nervous system neurotransmitters and plays a crucial role in integrating the body's overall response to stress.

Administration of CRF directly to the brain elicits behavioral, physiological and endocrine responses identical to those observed for an animal exposed to a stressful environment.

Accordingly, clinical data suggests that CRF receptor antagonists may represent novel antidepressant and/or anxiolytic drugs that may be useful in the treatment of the neuropsychiatric disorders manifesting hypersecretion of CRF.

The first CRF receptor antagonists were peptides (see, e.g., Rivier et al., U.S. Patent No. 4,605,642; Rivier et al., Science 224: 889, 1984). While these peptides established that CRF receptor antagonists can attenuate the pharmacological responses to CRF, peptide CRF receptor antagonists suffer from the usual drawbacks of peptide therapeutics including lack of stability and limited oral activity. More recently, small molecule CRF receptor antagonists have been reported.


WO 95/10506 describes inter alia compounds of general formula (A) with general CRF antagonist activity



wherein Y may be CR<sub>20</sub>; V may be nitrogen, Z may be carbon, R<sub>3</sub> may correspond to an amine derivative and R<sub>4</sub> may be taken together with R<sub>20</sub> to form a 5-membered ring and is -


CH(R<sub>29</sub>) when R<sub>29</sub> is -CH(R<sub>30</sub>). There are no specific disclosures of compounds corresponding to this definition.

WO 95/33750 (and in an analogous way WO01/53263 and EP773023) describes compounds of general formula (B) having CRF antagonistic activity,



in which A and Y may be nitrogen and carbon and B may correspond to an amine derivative. There are no specific disclosures of compounds corresponding to this definition.

The present invention constitutes a novel selection of the invention object of an International patent application filed by the same Applicant WO 03/008412, describing compounds of the following general formula C:



and including in the specification the preparation of the following compounds:

1-(2,4-bis-trifluoromethyl-phenyl)-6-methyl-4-(3-thiazol-2-yl-pyrazol-1-yl)-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine;

3-methyl-4-[6-methyl-4-(3-thiazol-2-yl-pyrazol-1-yl)-2,3-dihydro-1H-pyrrolo[2,3-b]pyridin-1-yl]-benzonitrile;

4-[6-methyl-4-(3-thiazol-2-yl-pyrazol-1-yl)-2,3-dihydro-1H-pyrrolo[2,3-b]pyridin-1-yl]-3-trifluoromethyl-benzonitrile;

6-methyl-1-(2-methyl-4-trifluoromethoxy-phenyl)-4-(3-thiazol-2-yl-pyrazol-1-yl)-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine;

1-(4-methoxy-2-methyl-phenyl)-6-methyl-4-(3-thiazol-2-yl-pyrazol-1-yl)-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine

1-(2,4-bis-trifluoromethyl-phenyl)-6-methyl-4-(3-pyridin-2-yl-pyrazol-1-yl)-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine


4-[1,3']bipyrazolyl-1'-yl-1-(2,4-bis-trifluoromethyl-phenyl)-6-methyl-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine.

Due to the physiological significance of CRF, the development of biologically-active small molecules having significant CRF receptor binding activity and which are capable of antagonizing the CRF receptor remains a desirable goal. Such CRF receptor antagonists would be useful in the treatment of endocrine, psychiatric and neurologic conditions or illnesses, including stress-related disorders in general.

While significant strides have been made toward achieving CRF regulation through administration of CRF receptor antagonists, there remains a need in the art for effective small molecule CRF receptor antagonists. There is also a need for pharmaceutical compositions containing such CRF receptor antagonists, as well as methods relating to the use thereof to treat, for example, stress-related disorders. The present invention fulfills these needs, and provides other related advantages.

In particular the invention relates to novel compounds which are potent and specific antagonists of corticotropin-releasing factor (CRF) receptors.

The present invention provides compounds of formula (I) including stereoisomers, prodrugs and pharmaceutically acceptable salts or solvates thereof



wherein

- R is aryl or heteroaryl, each of which may be substituted by 1 or more Z groups;
- R<sub>1</sub> is hydrogen, C3-C7 cycloalkyl, C1-C6 alkyl, C1-C6 alkoxy, C1-C6 thioalkyl, C2-C6 alkenyl, C2-C6 alkynyl, halo C1-C6 alkyl, halo C1-C6 alkoxy, halogen, NR<sub>6</sub>R<sub>7</sub> or cyano;
- R<sub>2</sub> and R<sub>3</sub> together with N form a 5-6 membered aromatic heterocycle, which is substituted by at least one group R<sub>8</sub> and may be further substituted by 1 to 3 R<sub>9</sub> groups;
- R<sub>4</sub> hydrogen, C1-C6 alkyl, halogen or halo C1-C6 alkyl;
- R<sub>5</sub> is a C1-C4 alkyl, -OR<sub>6</sub> or -NR<sub>6</sub>R<sub>7</sub>;
- R<sub>6</sub> is hydrogen or C1-C6 alkyl;
- R<sub>7</sub> is hydrogen or C1-C6 alkyl;
- R<sub>8</sub> is a 5-6 membered aromatic heterocycle, which may be substituted by 1 to 4 R<sub>10</sub> groups;
- R<sub>9</sub> is hydrogen, C3-C7 cycloalkyl, C3-C7 cycloalkenyl, C1-C6 alkyl, C1-C6 alkoxy, halo C1-C6 alkyl, halo C1-C6 alkoxy, hydroxy, halogen, nitro, cyano, -C(O)R<sub>6</sub>, C(O)NR<sub>6</sub>R<sub>7</sub>, phenyl which may be substituted by 1 to 4 R<sub>10</sub> groups;
- R<sub>10</sub> is C1-C6 alkyl, halo C1-C2 alkyl, halogen, nitro, hydroxy, halo C1-C6 alkoxy, C1-C6 alkoxy, or cyano;

Z is selected in a group consisting from halogen, C1-C6 alkyl, C1-C6 alkoxy, halo C1-C6 alkyl, C2-C6 alkenyl, C2-C6 alkynyl, halo C1-C6 alkoxy, -C(O)R<sub>6</sub>, -NR<sub>6</sub>R<sub>7</sub>, nitro, cyano, and a group R<sub>8</sub>;  
 and when R<sub>4</sub> is hydrogen, R<sub>2</sub> and R<sub>3</sub> together with N form a pyrazolyl group substituted with at least one R<sub>8</sub> corresponding to a thiazolyl group, and R corresponds to a phenyl group, and the substituent Z is at least one nitro group,  
 then in the compounds of formula (I) the nitro group is not present in the ortho position with respect to the nitrogen atom present in the 5-membered ring, named as B;

and the compounds of formula (I) don't include the following:

1-(2,4-bis-trifluoromethyl-phenyl)-6-methyl-4-(3-thiazol-2-yl-pyrazol-1-yl)-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine;  
 3-methyl-4-[6-methyl-4-(3-thiazol-2-yl-pyrazol-1-yl)-2,3-dihydro-1H-pyrrolo[2,3-b]pyridin-1-yl]-benzonitrile;  
 4-[6-methyl-4-(3-thiazol-2-yl-pyrazol-1-yl)-2,3-dihydro-1H-pyrrolo[2,3-b]pyridin-1-yl]-3-trifluoromethyl-benzonitrile;  
 6-methyl-1-(2-methyl-4-trifluoromethoxy-phenyl)-4-(3-thiazol-2-yl-pyrazol-1-yl)-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine;  
 1-(4-methoxy-2-methyl-phenyl)-6-methyl-4-(3-thiazol-2-yl-pyrazol-1-yl)-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine  
 1-(2,4-bis-trifluoromethyl-phenyl)-6-methyl-4-(3-pyridin-2-yl-pyrazol-1-yl)-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine  
 4-[1,3]bipyrazolyl-1'-yl-1-(2,4-bis-trifluoromethyl-phenyl)-6-methyl-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine.

The compounds of the present invention may be in the form of and/or may be administered as a pharmaceutically acceptable salt. For a review on suitable salts see Berge et al, J. Pharm. Sci., 1977, 66, 1-19.

Typically, a pharmaceutical acceptable salt may be readily prepared by using a desired acid or base as appropriate. The salt may precipitate from solution and be collected by filtration or may be recovered by evaporation of the solvent.

Suitable addition salts are formed from acids which form non-toxic salts and examples are hydrochloride, hydrobromide, hydriodide, sulphate, bisulphate, nitrate, phosphate, hydrogen phosphate, acetate, maleate, malate, fumarate, lactate, tartrate, citrate, formate, gluconate, succinate, piruvate, oxalate, oxaloacetate, trifluoroacetate, saccharate, benzoate, methansulphonate, ethanesulphonate, benzenesulphonate, p-toluensulphonate, methanesulphonic, ethanesulphonic, p-toluenesulphonic, and isethionate.

Pharmaceutically acceptable base salts include ammonium salts, alkali metal salts such as those of sodium and potassium, alkaline earth metal salts such as those of calcium and magnesium and salts with organic bases, including salts of primary, secondary and tertiary amines, such as isopropylamine, diethylamine, ethanolamine, trimethylamine, dicyclohexylamine and N-methyl-D-glucamine.

Those skilled in the art of organic chemistry will appreciate that many organic compounds can form complexes with solvents in which they are reacted or from which they are precipitated or crystallized. These complexes are known as "solvates". For example, a complex with water is known as a "hydrate". Solvates of the compound of the invention are within the scope of the invention.

In addition, prodrugs are also included within the context of this invention.

As used herein, the term "prodrug" means a compound which is converted within the body, e.g. by hydrolysis in the blood, into its active form that has medical effects. Pharmaceutically acceptable prodrugs are described in T. Higuchi and V. Stella, Prodrugs as Novel Delivery Systems, Vol. 14 of the A.C.S. Symposium Series, Edward B. Roche, ed., Bioreversible Carriers in Drug Design, American Pharmaceutical Association and Pergamon Press, 1987, and in D. Fleisher, S. Ramon and H. Barbra "Improved oral drug delivery: solubility limitations overcome by the use of prodrugs", Advanced Drug Delivery Reviews (1996) 19(2) 115-130, each of which are incorporated herein by reference.

Prodrugs are any covalently bonded carriers that release a compound of structure (I) *in vivo* when such prodrug is administered to a patient. Prodrugs are generally prepared by modifying functional groups in a way such that the modification is cleaved, either by routine manipulation or *in vivo*, yielding the parent compound. Prodrugs include, for example, compounds of this invention wherein hydroxy, amine or sulfhydryl groups are bonded to any group that, when administered to a patient, cleaves to form the hydroxy, amine or sulfhydryl groups. Thus, representative examples of prodrugs include (but are not limited to) acetate, formate and benzoate derivatives of alcohol, sulfhydryl and amine functional groups of the compounds of structure (I). Further, in the case of a carboxylic acid (-COOH), esters may be employed, such as methyl esters, ethyl esters, and the like. Esters may be active in their own right and/or be hydrolysable under *in vivo* conditions in the human body. Suitable pharmaceutically acceptable *in vivo* hydrolysable ester groups include those which break down readily in the human body to leave the parent acid or its salt.

With regard to stereoisomers, the compounds of structure (I) may have one or more asymmetric carbon atom and may occur as racemates, racemic mixtures and as individual enantiomers or diastereomers. All such isomeric forms are included within the present invention, including mixtures thereof.

Where a compound of the invention contains an alkenyl or alkenylene group, *cis* (E) and *trans* (Z) isomerism may also occur. The present invention includes the individual

stereoisomers of the compound of the invention and, where appropriate, the individual tautomeric forms thereof, together with mixtures thereof.

Separation of diastereoisomers or cis and trans isomers may be achieved by conventional techniques, e.g. by fractional crystallisation, chromatography or H.P.L.C. of a stereoisomeric mixture of the agent may also be prepared from a corresponding optically pure intermediate or by resolution, such as H.P.L.C. of the corresponding racemate using a suitable chiral support or by fractional crystallisation of the diastereoisomeric salts formed by reaction of the corresponding racemate with a suitable optically active acid or base, as appropriate.

Furthermore, some of the crystalline forms of the compounds of structure (I) may exist as polymorphs, which are included in the present invention.

The term C1-C6 alkyl as used herein as a group or a part of the group refers to a linear or branched alkyl group containing from 1 to 6 carbon atoms; examples of such groups include methyl, ethyl, propyl, isopropyl, n-butyl, isobutyl, tert butyl, pentyl or hexyl.

The term C3-C7 cycloalkyl group means a saturated hydrocarbon ring of 3 to 7 carbon atom such as, for example, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl or cycloheptyl; while unsaturated cycloalkyls include cyclopentenyl and cyclohexenyl, and the like.

The term C3-C7 cycloalkenyl group means a non aromatic monocyclic hydrocarbon ring of 3 to 7 carbon atom, containing from one to two unsaturation such as, cyclopentenyl and cyclohexenyl, and the like.

The term halogen refers to a fluorine, chlorine, bromine or iodine atom.

The term halo C1-C6 alkyl, or halo C1-C2 alkyl means an alkyl group having one or more carbon atoms and wherein at least one hydrogen atom is replaced with halogen such as for example a trifluoromethyl group and the like.

The term C2-C6 alkenyl defines straight or branched chain hydrocarbon radicals containing one or more double bond and having from 2 to 6 carbon atoms such as, for example, ethenyl, 2-propenyl, 3-butenyl, 2-butenyl, 2-pentenyl, 3-pentenyl, 3-methyl-2-butenyl or 3-hexenyl and the like.

The term C1-C6 alkoxy group may be a linear or a branched chain alkoxy group, for example methoxy, ethoxy, propoxy, prop-2-oxy, butoxy, but-2-oxy or methylprop-2-oxy and the like.

The term C1-C6 thioalkyl may be a linear or a branched chain thioalkyl group, for example thiomethyl, thioethyl, thiopropyl, thioisopropyl, thiobutyl, thiosec-butyl, thiotert-butyl and the like.

The term halo C1-C6 alkoxy group may be a C1-C6 alkoxy group as defined before substituted with at least one halogen, preferably fluorine, such as OCHF<sub>2</sub>, or OCF<sub>3</sub>.

The term C2-C6 alkynyl defines straight or branched chain hydrocarbon radicals containing one or more triple bond and having from 2 to 6 carbon atoms including acetylenyl, propynyl, 1-butynyl, 1-pentynyl, 3-methyl-1-butynyl and the like.

The term aryl means an aromatic carbocyclic moiety such as phenyl, biphenyl or naphthyl.


The term heteroaryl means an aromatic heterocycle ring of 5 to 10 members and having at least one heteroatom selected from nitrogen, oxygen and sulfur, and containing at least 1 carbon atom, including both mono- and bicyclic ring systems.

Representative heteroaryls include (but are not limited to) furyl, benzofuranyl, thiophenyl, benzothiophenyl, pyrrolyl, indolyl, isoindolyl, azaindolyl, pyridyl, quinoliny, isoquinoliny, oxazolyl, isooxazolyl, benzoxazolyl, pyrazolyl, imidazolyl, benzimidazolyl, thiazolyl, benzothiazolyl, isothiazolyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl, cinnoliny, phthalazinyl, triazolyl, tetrazolyl, benzopyrazolyl, and quinazolinyl.

The term 5-6 membered aromatic heterocycle means, a 5-6 monocyclic heterocyclic ring which is aromatic, and which contains from 1 to 4 heteroatoms independently selected from nitrogen, oxygen and sulfur, and wherein the nitrogen and sulfur heteroatoms may be optionally oxidized, and the nitrogen heteroatom may be optionally quaternized. Heterocycles include some of the heteroaryls as defined above. The heterocycle may be attached via any heteroatom or carbon atom. Thus, the term include (but are not limited to) furyl, thiophenyl, pyrrolyl, tetrazolyl, pyrimidinyl, pyridinyl, pyrazinyl, pyrazolyl, thiazolyl, triazolyl, imidazolyl, oxadiazolyl, oxazolyl, and the like. Since such 5-6 membered aromatic heterocycles may be substituted by hydroxy groups, the possible tautomeric forms are also part of the present invention.

Representative compounds of this invention include compounds of general formula (II), corresponding to compounds of formula (I) in which R<sub>4</sub> is hydrogen and R, R<sub>1</sub>, R<sub>2</sub> and R<sub>3</sub> are defined as above.






and when  $R_4$  is hydrogen,  $R_2$  and  $R_3$  together with N form a pyrazolyl group substituted with at least one  $R_6$  corresponding to a thiazolyl group, and R corresponds to a phenyl group, and the substituent Z is at least one nitro group,

then In the compounds of formula (II) the nitro group is not present in the ortho position with respect to the nitrogen atom present in the 5-membered ring, named as B;

and the compounds of formula (II) don't include the following:

- 1-(2,4-bis-trifluoromethyl-phenyl)-6-methyl-4-(3-thiazol-2-yl-pyrazol-1-yl)-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine;
- 3-methyl-4-[6-methyl-4-(3-thiazol-2-yl-pyrazol-1-yl)-2,3-dihydro-1H-pyrrolo[2,3-b]pyridin-1-yl]-benzonitrile;
- 4-[6-methyl-4-(3-thiazol-2-yl-pyrazol-1-yl)-2,3-dihydro-1H-pyrrolo[2,3-b]pyridin-1-yl]-3-trifluoromethyl-benzonitrile;
- 6-methyl-1-(2-methyl-4-trifluoromethoxy-phenyl)-4-(3-thiazol-2-yl-pyrazol-1-yl)-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine;
- 1-(4-methoxy-2-methyl-phenyl)-6-methyl-4-(3-thiazol-2-yl-pyrazol-1-yl)-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine
- 1-(2,4-bis-trifluoromethyl-phenyl)-6-methyl-4-(3-pyridin-2-yl-pyrazol-1-yl)-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine
- 4-[1,3']bipyrazolyl-1'-yl-1-(2,4-bis-trifluoromethyl-phenyl)-6-methyl-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine.

In particular compounds of formula (III) are preferred, corresponding to the compounds of formula (II)



in which

$R_2$  and  $R_3$  together with N form a pyrazolyl group;

n is an integer from 1 to 2;


and R,  $R_1$ ,  $R_6$  and  $R_9$  have the meanings as defined before;

and when  $R_4$  is hydrogen,  $R_2$  and  $R_3$  together with N form a pyrazolyl group substituted with at least one  $R_8$  corresponding to a thiazolyl group, and R corresponds to a phenyl group, and the substituent Z is at least one nitro group,  
 then in the compounds of formula (III) the nitro group is not present in the ortho position with respect to the nitrogen atom present in the 5-membered ring, named as B;

and the compounds of formula (III) don't include the following:

1-(2,4-bis-trifluoromethyl-phenyl)-6-methyl-4-(3-thiazol-2-yl-pyrazol-1-yl)-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine;  
 3-methyl-4-[6-methyl-4-(3-thiazol-2-yl-pyrazol-1-yl)-2,3-dihydro-1H-pyrrolo[2,3-b]pyridin-1-yl]-benzonitrile;  
 4-[6-methyl-4-(3-thiazol-2-yl-pyrazol-1-yl)-2,3-dihydro-1H-pyrrolo[2,3-b]pyridin-1-yl]-3-trifluoromethyl-benzonitrile;  
 6-methyl-1-(2-methyl-4-trifluoromethoxy-phenyl)-4-(3-thiazol-2-yl-pyrazol-1-yl)-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine;  
 1-(4-methoxy-2-methyl-phenyl)-6-methyl-4-(3-thiazol-2-yl-pyrazol-1-yl)-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine  
 1-(2,4-bis-trifluoromethyl-phenyl)-6-methyl-4-(3-pyridin-2-yl-pyrazol-1-yl)-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine  
 4-[1,3]bipyrazolyl-1'-yl-1-(2,4-bis-trifluoromethyl-phenyl)-6-methyl-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine.


Preferred are the compounds of formula (IV), corresponding to compounds of formula (III),



in which:

- $R_8$  is a thiazolyl derivative;
- $n$  is an integer from 1 to 2;
- R corresponds to W and
- W is a pyridine derivative, which may be substituted by 1 to 4 Z groups, as defined above; and
- $R_1$ ,  $R_9$  and  $R_{10}$  are defined as above.


Particularly preferred are the compounds of formula (IVa).



in which W is defined as above.

Specific examples of compounds of general formula (IVa) are included in the Experimental Part.

Equally preferred are compounds of formula (V), corresponding to compounds of formula (III),



in which:

- $R_8$  is a thiazolyl derivative,;
- $n$  is an integer from 1 to 2;
- $R$  corresponds to  $W_1$  and
- $W_1$  is a phenyl derivative substituted by 2 to 4  $Z_1$  groups;
- $Z_1$  is selected in a group consisting from halogen, C1-C6 alkyl, C1-C6 alkoxy, halo C1-C6 alkyl, C2-C6 alkenyl, C2-C6 alkynyl, halo C1-C6 alkoxy,  $-C(O)R_6$ ,  $-NR_6R_7$ , cyano, and a group  $R_8$ ;


$R_1$ ,  $R_9$  and  $R_{10}$  are defined as above;

and the compounds of formula (V) don't include the following:

- 1-(2,4-bis-trifluoromethyl-phenyl)-6-methyl-4-(3-thiazol-2-yl-pyrazol-1-yl)-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine;
- 3-methyl-4-[6-methyl-4-(3-thiazol-2-yl-pyrazol-1-yl)-2,3-dihydro-1H-pyrrolo[2,3-b]pyridin-1-yl]-benzonitrile;
- 4-[6-methyl-4-(3-thiazol-2-yl-pyrazol-1-yl)-2,3-dihydro-1H-pyrrolo[2,3-b]pyridin-1-yl]-3-trifluoromethyl-benzonitrile;

6-methyl-1-(2-methyl-4-trifluoromethoxy-phenyl)-4-(3-thiazol-2-yl-pyrazol-1-yl)-2,3-dihydro-1*H*-pyrrolo[2,3-*b*]pyridine;  
 1-(4-methoxy-2-methyl-phenyl)-6-methyl-4-(3-thiazol-2-yl-pyrazol-1-yl)-2,3-dihydro-1*H*-pyrrolo[2,3-*b*]pyridine  
 1-(2,4-bis-trifluoromethyl-phenyl)-6-methyl-4-(3-pyridin-2-yl-pyrazol-1-yl)-2,3-dihydro-1*H*-pyrrolo[2,3-*b*]pyridine  
 4-[1,3']bipyrazolyl-1'-yl-1-(2,4-bis-trifluoromethyl-phenyl)-6-methyl-2,3-dihydro-1*H*-pyrrolo[2,3-*b*]pyridine.


Particularly preferred are the compounds of formula (Va),



in which  $W_1$  is defined as above.

Specific examples of compounds of general formula (Va) are included in the Experimental Part.

Equally preferred are compounds of formula (VI), corresponding to compounds of formula (III),




in which:

- $R_9$  is a thiazolyl derivative;
- $n$  is an integer from 1 to 2;
- $R$  corresponds to  $W_2$  and
- $W_2$  is a phenyl derivative, which may be substituted by 2 to 4  $Z$  groups as defined above provided that at least one  $Z$  group is nitro and further provided that the

nitro group is not in the ortho position with respect to the nitrogen atom of the 5-membered ring named as B; and  
 $R_1$ ,  $R_9$  and  $R_{10}$  are defined as above.

Even more preferred are the compounds of formula (V), in which R is a phenyl derivative containing a nitro group in para position respect to the nitrogen atom of the 5-membered ring B.


Particularly preferred are the compounds of formula (VIa),



in which  $W_2$  is defined as above.

Specific examples of compounds of general formula (VIa) are included in the Experimental Part.

Equally preferred are compounds of formula (VII),




in which

$R_8$  is a pyrazolyl derivative;

$n$  is an integer from 1 to 2;

$R$ ,  $R_9$  and  $R_{10}$  are defined as above.


Particularly preferred are the compounds of formula (VIIa),



in which  $W_1$  is defined as above.

Specific examples of compounds of general formula (VIIa) are included in the Experimental Part.


Equally preferred are the compounds of formula (VIII),



in which

- $R_9$  is a pyridine derivative;
- $n$  is an integer from 1 to 2;
- $m$  is an integer from 1 to 3;
- $R$ ,  $R_9$  and  $R_{10}$  are defined as above.

Particularly preferred are the compounds of formula (VIIIa), corresponding to the compounds of formula (VIII)



in which  $W_1$  is defined as above.

Specific examples of compounds of general formula (VIIIa) are included in the Experimental Part.

Preferred compounds according to the invention are:

6-methyl-1-[6-(methyloxy)-2-(trifluoromethyl)-3-pyridinyl]-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine;  
1-[2,6-bis(methyloxy)-3-pyridinyl]-6-methyl-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine;  
N,N,4-trimethyl-5-[6-methyl-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-2,3-dihydro-1H-pyrrolo[2,3-b]pyridin-1-yl]-2-pyridinamine;  
1-(2-difluoromethyl-4-methoxy-phenyl)-6-methyl-4-(3-thiazol-2-yl-pyrazol-1-yl)-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine;  
1-(2-chloro-4-methoxy-phenyl)-6-methyl-4-(3-thiazol-2-yl-pyrazol-1-yl)-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine;  
1-[2,4-bis(methyloxy)phenyl]-6-methyl-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine;  
3-chloro-4-[6-methyl-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-2,3-dihydro-1H-pyrrolo[2,3-b]pyridin-1-yl]benzonitrile;  
4-[6-methyl-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-2,3-dihydro-1H-pyrrolo[2,3-b]pyridin-1-yl]-3-[(trifluoromethyl)oxy]benzonitrile;  
3-ethyl-4-[6-methyl-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-2,3-dihydro-1H-pyrrolo[2,3-b]pyridin-1-yl]benzonitrile;  
1-(4-fluoro-2-methylphenyl)-6-methyl-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine;  
6-methyl-1-[2-methyl-4-(1H-pyrazol-1-yl)phenyl]-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine;  
6-methyl-1-[4-nitro-2-(trifluoromethyl)phenyl]-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine;  
4-[4-(1H-1,3'-bipyrazol-1'-yl)-6-methyl-2,3-dihydro-1H-pyrrolo[2,3-b]pyridin-1-yl]-3-methylbenzonitrile;  
4-[4-(1H-1,3'-bipyrazol-1'-yl)-6-methyl-2,3-dihydro-1H-pyrrolo[2,3-b]pyridin-1-yl]-3-(trifluoromethyl)benzonitrile;  
4-[4-(1H-1,3'-bipyrazol-1'-yl)-6-methyl-2,3-dihydro-1H-pyrrolo[2,3-b]pyridin-1-yl]-3-chlorobenzonitrile;  
1'-[6-methyl-1-[2-methyl-4-(methyloxy)phenyl]-2,3-dihydro-1H-pyrrolo[2,3-b]pyridin-4-yl]-1H-1,3'-bipyrazole;  
3-methyl-4-[6-methyl-4-[3-(2-pyridinyl)-1H-pyrazol-1-yl]-2,3-dihydro-1H-pyrrolo[2,3-b]pyridin-1-yl]benzonitrile;  
3-chloro-4-[6-methyl-4-[3-(2-pyridinyl)-1H-pyrazol-1-yl]-2,3-dihydro-1H-pyrrolo[2,3-b]pyridin-1-yl]benzonitrile;


4-{6-methyl-4-[3-(2-pyridinyl)-1*H*-pyrazol-1-yl]-2,3-dihydro-1*H*-pyrrolo[2,3-*b*]pyridin-1-yl}-3-(trifluoromethyl)benzonitrile.

In general, the compounds of structure (I) may be made according to the organic synthesis techniques known to those skilled in this field, as well as by the representative methods set forth in the Examples.

Compounds of formula (I), and salts and solvates thereof, may be prepared by the general methods outlined hereinafter. In the following description, the groups R, R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub>, R<sub>5</sub>, R<sub>6</sub>, R<sub>7</sub>, R<sub>8</sub>, R<sub>9</sub>, R<sub>10</sub>, Z, W, W<sub>1</sub>, W<sub>2</sub>, m and n have the meanings as previously defined for compounds of formula (I) unless otherwise stated.

Compounds of formula (I) may be conveniently prepared, starting from compounds of formula (IIIB), according to the following Scheme 1:

Scheme 1




in which



- step a stands for conversion of the leaving group L, selected in a group consisting from: halogen or reactive residue of sulphonic acid (e.g. mesylate, tosylate), preferably chloride, in the amino group of compounds (IVB), by reaction with the suitable amine  $\text{NR}_2\text{R}_3$  in basic conditions;
- step b stands for reduction of the ester group with a suitable reducing agent (such as DIBAL-H) to hydroxy group of compounds (VB);
- step c stands for oxidation of the hydroxy group with a suitable oxidising agent (such as Dess-Martin periodinane) to aldehyde group of compound (VIB);
- step d stands for formation of the aldehyde group of compounds (VIII B) by Wittig reaction in the usual conditions, through formation of enol ether followed by acid hydrolysis (step e);
- step f stands for reduction of the aldehyde group with a suitable reducing agent (such as  $\text{NaBH}_4$ ) to hydroxy group of compounds (IXB);
- step g stands for conversion of the hydroxy group in the suitably protected hydroxy group of compounds (XB)(such as TBS: *tert*-butyldimethylsilyl);
- step h stands for Buchwald reaction by coupling with the suitable amine  $\text{RNH}_2$ ;
- step i stands for deprotection reaction to give the hydroxy group of compounds (XIIB);
- step l stands for intramolecular cyclisation after conversion of the hydroxy group of compounds (XIIB) in a suitable leaving group (such as bromide, by reaction with  $\text{CBr}_4$  and  $\text{PPh}_3$ ) to give the final compounds (I).

Alternatively, compounds of formula (I) may be conveniently prepared according to the following Scheme 2:

Scheme 2




in which

- step a' stands for conversion of the hydroxy group in a suitable leaving group L' of compounds (XIIIB), which, independently from L, has the same definition (e.g. mesylate, by reaction with MsCl and Et<sub>3</sub>N);
- step b' stands for conversion of L' in the cyano derivative of compounds (XIVB) by reaction with, e.g. KCN in an aprotic dipolar solvent, like DMF;
- step c' stands for reduction of the cyano group with a suitable reducing agent (such as BH<sub>3</sub>-THF) to the amino group of compound (XVB);
- step d' stands for intramolecular cyclisation of compounds (XVB) by heating in a suitable solvent (such as NMP) at high temperature or intramolecular Copper catalysed cyclisation;
- step e' see step h above. The reaction is done with a suitable aryl halide to give the final compounds (I).

Alternatively, compounds of formula (I) may be conveniently prepared according to the following Scheme 3:

Scheme 3



in which:


- step a'' stands for the formation of the pyrrolidinone moiety of compounds (XVIIIIB), which will form the cycle B present in the final compounds (I), by reacting the compounds (XVIIIB) with a reactive derivative of the butyric acid, such as 4-chlorobutyl chloride; followed by a cyclisation reaction in basic conditions (e.g. KOtBu);
- step b'' stands for amidine formation by reacting the compounds (XIXB) with 3-aminocrotonate and POCl<sub>3</sub>;
- step c'' stands for the cyclisation of the compounds (XIXB) in basic conditions (e.g. NaH) to give the pyridinone precursor of cycle A in the final compounds (I);
- step d'' stands for the formation of a reactive derivative (i.e. a leaving group, Lg) of the pyridinone (for example selected in a group consisting by triflate,

- halogen, mesylate) of compounds (XXIB) by reaction with, for example, triflic anhydride;
- step e'' stands for nucleophilic displacement of the leaving group of compounds (XXIB) to give the iodinated compounds (XXIIB);
- step f'' stands for the arylation reaction with the suitable pyrazole derivative by a metal catalysed coupling reaction (for example a Buchwald reaction) procedure to give the final compounds (I).

Another aspect of the present invention is to furnish a convenient process for preparing derivatives in which the group R contains  $-\text{CHF}_2$  as one or more of the groups Z.

Such process comprises the following steps described in Scheme 4. The process has been exemplified for R corresponding to phenyl derivative, but it is not limiting its appropriateness to the various meanings of R, as defined above.

Scheme 5



- step a''' stands for the reduction of the acid group present in (XXIIIB) to the hydroxy group of compounds (XXIVB) by a suitable reducing agent (such as treatment with cyanuric chloride and NMM and then reduction with  $\text{NaBH}_4$ );
- step b''' stands for the oxidation to the aldehyde group of compounds (XXVB) by a suitable oxidising agent (such as Dess Martin periodinane);
- step c''' stands for the fluorination of the aldehyde group by a suitable fluorinating agent (such as DAST: (diethylamino)sulphurtrifluoride);
- step d''' stands for the final reduction of the nitro group to the amino group of compounds (XXVIIIB) by a suitable reducing agent.

Compounds of formula (IIIB), (VB), (XVIIB) and (XXIIIB) are known compounds or may be prepared according to known methods in the literature.

Compounds of formula (III), (IIIa), (IV), (IVa), (V), (Va), (VI), (VIa), (VII), (VIIa), (VIII), and (VIIIa) may be prepared according to the previous Schemes 1, 2 and 3, once prepared the heterocyclic reactive residue according to known methods to the skilled in the art.

Those skilled in the art will appreciate that in the preparation of the compound of the invention or a solvate thereof it may be necessary and/or desirable to protect one or more sensitive groups in the molecule to prevent undesirable side reactions. Suitable protecting groups for use according to the present invention are well known to those skilled in the art and may be used in a conventional manner. See, for example, "Protective groups in organic synthesis" by T.W. Greene and P.G.M. Wuts (John Wiley & sons 1991) or "Protecting Groups" by P.J. Kocienski (Georg Thieme Verlag 1994). Examples of suitable amino protecting groups include acyl type protecting groups (e.g. formyl, trifluoroacetyl, acetyl), aromatic urethane type protecting groups (e.g. benzyloxycarbonyl (Cbz) and substituted Cbz), aliphatic urethane protecting groups (e.g. 9-fluorenylmethoxycarbonyl (Fmoc), t-butyloxycarbonyl (Boc), isopropyloxycarbonyl, cyclohexyloxycarbonyl) and alkyl type protecting groups (e.g. benzyl, trityl, chlorotriyl). Examples of suitable oxygen protecting groups may include for example alkyl silyl groups, such as trimethylsilyl or tert-butyldimethylsilyl; alkyl ethers such as tetrahydropyranyl or tert-butyl; or esters such as acetate

Pharmaceutical acceptable salts may also be prepared from other salts, including other pharmaceutically acceptable salts, of the compound of formula (I) using conventional methods.

The compounds of formula (I) may readily be isolated in association with solvent molecules by crystallisation or evaporation of an appropriate solvent to give the corresponding solvates.

When a specific enantiomer of a compound of general formula (I) is required, this may be obtained for example by resolution of a corresponding enantiomeric mixture of a compound of formula (I) using conventional methods. Thus the required enantiomer may be obtained from the racemic compound of formula (I) by use of chiral HPLC procedure.

The subject invention also includes isotopically-labelled compounds, which are identical to those recited in formula (I) and following, but for the fact that one or more atoms are replaced by an atom having an atomic mass or mass number different from the atomic mass or mass number usually found in nature. Examples of isotopes that can be incorporated into compounds of the invention and pharmaceutically acceptable salts thereof include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorous, sulphur, fluorine, iodine, and chlorine, such as  $^2\text{H}$ ,  $^3\text{H}$ ,  $^{11}\text{C}$ ,  $^{13}\text{C}$ ,  $^{14}\text{C}$ ,  $^{15}\text{N}$ ,  $^{17}\text{O}$ ,  $^{18}\text{O}$ ,  $^{31}\text{P}$ ,  $^{32}\text{P}$ ,  $^{35}\text{S}$ ,  $^{18}\text{F}$ ,  $^{38}\text{Cl}$ ,  $^{123}\text{I}$  and  $^{125}\text{I}$ .

Compounds of the present invention and pharmaceutically acceptable salts of said compounds that contain the aforementioned isotopes and/or other isotopes of other atoms are within the scope of the present invention. Isotopically-labelled compounds of the present invention, for example those into which radioactive isotopes such as  $^3\text{H}$ ,  $^{14}\text{C}$  are incorporated, are useful in drug and/or substrate tissue distribution assays. Tritiated, i.e.,  $^3\text{H}$ , and carbon-14, i.e.,  $^{14}\text{C}$ , isotopes are particularly preferred for their ease of preparation

and detectability.  $^{11}\text{C}$  and  $^{18}\text{F}$  isotopes are particularly useful in PET (positron emission tomography), and  $^{125}\text{I}$  isotopes are particularly useful in SPECT (single photon emission computerized tomography), all useful in brain imaging. Further, substitution with heavier isotopes such as deuterium, i.e.,  $^2\text{H}$ , can afford certain therapeutic advantages resulting from greater metabolic stability, for example increased *in vivo* half-life or reduced dosage requirements and, hence, may be preferred in some circumstances. Isotopically labelled compounds of formula I and following of this invention can generally be prepared by carrying out the procedures disclosed in the Schemes and/or in the Examples below, by substituting a readily available isotopically labelled reagent for a non-isotopically labelled reagent.

The CRF receptor antagonists of the present invention demonstrate activity at the CRF receptor site including CRF 1 and CRF 2 receptors and may be used in the treatment of conditions mediated by CRF or CRF receptors.

The effectiveness of a compound as a CRF receptor antagonist may be determined by various assay methods. Suitable CRF antagonists of this invention are capable of inhibiting the specific binding of CRF to its receptor and antagonizing activities associated with CRF. A compound of structure (I) may be assessed for activity as a CRF antagonist by one or more generally accepted assays for this purpose, including (but not limited to) the assays disclosed by DeSouza et al. (J. Neuroscience 7: 88,1987) and Battaglia et al. (Synapse 1: 572,1987).

The CRF receptors-binding assay was performed by using the homogeneous technique of scintillation proximity (SPA). The ligand binds to recombinant membrane preparation expressing the CRF receptors which in turn bind to wheatgerm agglutinin-coated SPA beads. In the Experimental Part will be disclosed the details of the experiments.

With reference to CRF receptor binding affinities, CRF receptor antagonists of this invention have a  $K_i$  less than  $10\ \mu\text{M}$ . In a preferred embodiment of this invention, a CRF receptor antagonist has a  $K_i$  comprised in a range from  $0.1\ \mu\text{M}$  and  $10\ \mu\text{M}$ .

In a more preferred embodiment the value of  $K_i$  is less than  $0.1\ \mu\text{M}$ . As set forth in greater detail below, the  $K_i$  values of representative compounds of this invention were assayed by the methods set forth in Example 6.

Compounds of the invention are useful in the treatment of central nervous system disorders where CRF receptors are involved. In particular in the treatment or prevention of major depressive disorders including bipolar depression, unipolar depression, single or recurrent major depressive episodes with or without psychotic features, catatonic features, melancholic features, atypical features or postpartum onset, the treatment of anxiety and the treatment of panic disorders. The term anxiety includes anxiety disorders, such as panic disorders with or without agoraphobia, agoraphobia, phobias, for example, social phobias or

agoraphobia, obsessive-compulsive disorder, stress disorders including post-traumatic stress disorders, generalised anxiety disorders, acute stress disorders and mixed anxiety-depression disorders. Other mood disorders encompassed within the term major depressive disorders include dysthymic disorder with early or late onset and with or without atypical features, neurotic depression, post traumatic stress disorders and social phobia; dementia of the Alzheimer's type, with early or late onset, with depressed mood; vascular dementia with depressed mood; mood disorders induced by alcohol, amphetamines, cocaine, hallucinogens, inhalants, opioids, phencyclidine, sedatives, hypnotics, anxiolytics and other substances; schizoaffective disorder of the depressed type; and adjustment disorder with depressed mood. Major depressive disorders may also result from a general medical condition including, but not limited to, myocardial infarction, diabetes, miscarriage or abortion, etc.

Compounds of the invention are also useful in the treatment or prevention of schizophrenic disorders including paranoid schizophrenia, disorganised schizophrenia, catatonic schizophrenia, undifferentiated schizophrenia, residual schizophrenia.

Compounds of the invention are useful as analgesics. In particular they are useful in the treatment of traumatic pain such as postoperative pain; traumatic avulsion pain such as brachial plexus; chronic pain such as arthritic pain such as occurring in osteo-, rheumatoid or psoriatic arthritis; neuropathic pain such as post-herpetic neuralgia, trigeminal neuralgia, segmental or intercostal neuralgia, fibromyalgia, causalgia, peripheral neuropathy, diabetic neuropathy, chemotherapy-induced neuropathy, AIDS related neuropathy, occipital neuralgia, geniculate neuralgia, glossopharyngeal neuralgia, reflex sympathetic dystrophy, phantom limb pain; various forms of headache such as migraine, acute or chronic tension headache, temporomandibular pain, maxillary sinus pain, cluster headache; odontalgia; cancer pain; pain of visceral origin; gastrointestinal pain; nerve entrapment pain; sport's injury pain; dysmenorrhoea; menstrual pain; meningitis; arachnoiditis; musculoskeletal pain; low back pain e.g. spinal stenosis; prolapsed disc; sciatica; angina; ankylosing spondylitis; gout; burns; scar pain; itch; and thalamic pain such as post stroke thalamic pain.

Compounds of the invention are also useful for the treatment of dysfunction of appetite and food intake and in circumstances such as anorexia, anorexia nervosa and bulimia.

Compounds of the invention are also useful in the treatment of sleep disorders including dysomnia, insomnia, sleep apnea, narcolepsy, and circadian rhythmic disorders.

Compounds of the invention are also useful in the treatment or prevention of cognitive disorders. Cognitive disorders include dementia, amnesic disorders and cognitive disorders not otherwise specified.

Furthermore compounds of the invention are also useful as memory and/or cognition enhancers in healthy humans with no cognitive and/or memory deficit.

Compounds of the invention are also useful in the treatment of tolerance to and dependence on a number of substances. For example, they are useful in the treatment of dependence on nicotine, alcohol, caffeine, phencyclidine (phencyclidine like compounds), or in the treatment of tolerance to and dependence on opiates (e.g. cannabis, heroin, morphine) or benzodiazepines; in the treatment of cocaine, sedative hypnotic, amphetamine or amphetamine-related drugs (e.g. dextroamphetamine, methylamphetamine) addiction or a combination thereof.

Compounds of the invention are also useful as anti-inflammatory agents. In particular they are useful in the treatment of inflammation in asthma, influenza, chronic bronchitis and rheumatoid arthritis; in the treatment of inflammatory diseases of the gastrointestinal tract such as Crohn's disease, ulcerative colitis, postoperative gastric ileus (POI), inflammatory bowel disease (IBD) and non-steroidal anti-inflammatory drug induced damage; inflammatory diseases of the skin such as herpes and eczema; inflammatory diseases of the bladder such as cystitis and urge incontinence; and eye and dental inflammation.

Compounds of the invention are also useful in the treatment of allergic disorders, in particular allergic disorders of the skin such as urticaria, and allergic disorders of the airways such as rhinitis.

Compounds of the invention are also useful in the treatment of emesis, i.e. nausea, retching and vomiting. Emesis includes acute emesis, delayed emesis and anticipatory emesis. The compounds of the invention are useful in the treatment of emesis however induced. For example, emesis may be induced by drugs such as cancer chemotherapeutic agents such as alkylating agents, e.g. cyclophosphamide, carmustine, lomustine and chlorambucil; cytotoxic antibiotics, e.g. dactinomycin, doxorubicin, mitomycin-C and bleomycin; anti-metabolites, e.g. cytarabine, methotrexate and 5-fluorouracil; vinca alkaloids, e.g. etoposide, vinblastine and vincristine; and others such as cisplatin, dacarbazine, procarbazine and hydroxyurea; and combinations thereof; radiation sickness; radiation therapy, e.g. irradiation of the thorax or abdomen, such as in the treatment of cancer; poisons; toxins such as toxins caused by metabolic disorders or by infection, e.g. gastritis, or released during bacterial or viral gastrointestinal infection; pregnancy; vestibular disorders, such as motion sickness, vertigo, dizziness and Meniere's disease; post-operative sickness; gastrointestinal obstruction; reduced gastrointestinal motility; visceral pain, e.g. myocardial infarction or peritonitis; migraine; increased intracranial pressure; decreased intracranial pressure (e.g. altitude sickness); opioid analgesics, such as morphine; and gastro-oesophageal reflux disease, acid indigestion, over-indulgence of food or drink, acid stomach, sour stomach,

waterbrash/regurgitation, heartburn, such as episodic heartburn, nocturnal heartburn, and meal-induced heartburn and dyspepsia.

Compounds of the invention are of particular use in the treatment of gastrointestinal disorders such as irritable bowel syndrome (IBS); skin disorders such as psoriasis, pruritis and sunburn; vasospastic diseases such as angina, vascular headache and Reynaud's disease; cerebral ischaemia such as cerebral vasospasm following subarachnoid haemorrhage; fibrosing and collagen diseases such as scleroderma and eosinophilic fasciitis; disorders related to immune enhancement or suppression such as systemic lupus erythematosus and rheumatic diseases such as fibrositis; and cough.

Compounds of the invention are useful for the treatment of neurotoxic injury which follows cerebral stroke, thromboembolic stroke, hemorrhagic stroke, cerebral ischemia, cerebral vasospasm, hypoglycemia, hypoxia, anoxia, perinatal asphyxia cardiac arrest.

The invention therefore provides a compound of formula (I) or a pharmaceutically acceptable salt or solvate thereof for use in therapy, in particular in human medicine.

There is also provided as a further aspect of the invention the use of a compound of formula (I) or a pharmaceutically acceptable salt or solvate thereof in the preparation of a medicament for use in the treatment of conditions mediated by CRF.

In an alternative or further aspect there is provided a method for the treatment of a mammal, including man, in particular in the treatment of condition mediated by CRF, comprising administration of an effective amount of a compound of formula (I) or a pharmaceutically acceptable salt or a solvate thereof.

While it is possible that, for use in therapy, a compound of the present invention may be administered as the raw chemical, it is preferable to present the active ingredient as a pharmaceutical formulation e. g. when the agent is in admixture with a suitable pharmaceutical excipient, diluent or carrier selected with regard to the intended route of administration and standard pharmaceutical practice.

In a further aspect, the invention provides a pharmaceutical composition comprising at least one compound of the invention or a pharmaceutically acceptable derivative thereof in association with a pharmaceutically acceptable carrier and/or excipient. The carrier and/or excipient must be "acceptable" in the sense of being compatible with the other ingredients of the formulation and not deleterious to the recipient thereof.

Accordingly, the present invention further provides a pharmaceutical formulation comprising at least one compound of the invention or a pharmaceutically acceptable derivative thereof, in association with a pharmaceutically acceptable carrier and/or excipient. The carrier



and/or excipient must be "acceptable" in the sense of being compatible with the other ingredients of the formulation and not deleterious to the recipient thereof.

There is further provided by the present invention a process of preparing a pharmaceutical composition, which process comprises mixing at least one compound of the invention or a pharmaceutically acceptable derivative thereof, together with a pharmaceutically acceptable carrier and/or excipient.

The pharmaceutical compositions may be for human or animal usage in human and veterinary medicine and will typically comprise any one or more of a pharmaceutically acceptable diluent, carrier or excipient. Acceptable carriers or diluents for therapeutic use are well known in the pharmaceutical art, and are described, for example, in Remington's Pharmaceutical Sciences, Mack Publishing Co. (A. R. Gennaro edit. 1985). The choice of pharmaceutical carrier, excipient or diluent can be selected with regard to the intended route of administration and standard pharmaceutical practice. The pharmaceutical compositions may comprise as – or in addition to – the carrier, excipient or diluent any suitable binder(s), lubricant(s), suspending agent(s), coating agent(s), solubilising agent(s).

Preservatives, stabilisers, dyes and even flavouring agents may be provided in the pharmaceutical composition. Examples of preservatives include sodium benzoate, sorbic acid and esters of p-hydroxybenzoic acid. Antioxidants and suspending agents may be also used.

There may be different composition/formulation requirements dependent on the different delivery systems. By way of example, the pharmaceutical composition of the present invention may be formulated to be delivered using a mini-pump or by a mucosal route, for example, as a nasal spray or aerosol for inhalation or ingestible solution, or parenterally in which the composition is formulated by an injectable form, for delivery, by, for example, an intravenous, intramuscular or subcutaneous route. Alternatively, the formulation may be designed to be delivered by both routes.

Where the agent is to be delivered mucosally through the gastrointestinal mucosa, it should be able to remain stable during transit through the gastrointestinal tract; for example, it should be resistant to proteolytic degradation, stable at acid pH and resistant to the detergent effects of bile.

Where appropriate, the pharmaceutical compositions can be administered by inhalation, in the form of a suppository or pessary, topically in the form of a lotion, solution, cream, ointment or dusting powder, by use of a skin patch, orally in the form of tablets containing excipients such as starch or lactose, or in capsules or ovules either alone or in admixture with excipients, or in the form of elixirs, solutions or suspensions containing flavouring or colouring agents, or they can be injected parenterally, for example intravenously, intramuscularly or subcutaneously. For parenteral administration, the compositions may be

best used in the form of a sterile aqueous solution which may contain other substances, for example enough salts or monosaccharides to make the solution isotonic with blood. For buccal or sublingual administration the compositions may be administered in the form of tablets or lozenges which can be formulated in a conventional manner.

For some embodiments, the agents of the present invention may also be used in combination with a cyclodextrin. Cyclodextrins are known to form inclusion and non-inclusion complexes with drug molecules. Formation of a drugcyclodextrin complex may modify the solubility, dissolution rate, bioavailability and/or stability property of a drug molecule. Drug-cyclodextrin complexes are generally useful for most dosage forms and administration routes. As an alternative to direct complexation with the drug the cyclodextrin may be used as an auxiliary additive, e. g. as a carrier, diluent or solubiliser. Alpha-, betaand gamma-cyclodextrins are most commonly used and suitable examples are described in WO-A-91/11172, WO-A-94/02518 and WO-A-98/55148.

In a preferred embodiment, the agents of the present invention are delivered systemically (such as orally, buccally, sublingually), more preferably orally.

Hence, preferably the agent is in a form that is suitable for oral delivery.

It is to be understood that not all of the compounds need be administered by the same route. Likewise, if the composition comprises more than one active component, then those components may be administered by different routes.

The compounds of the invention may be milled using known milling procedures such as wet milling to obtain a particle size appropriate for tablet formation and for other formulation types. Finely divided (nanoparticulate) preparations of the compounds of the invention may be prepared by processes known in the art, for example see International Patent Application No. WO 02/00196 (SmithKline Beecham).

For oral administration, the pharmaceutical compositions may take the form of, for example, tablets or capsules prepared by conventional means with pharmaceutically acceptable excipients such as binding agents (e.g. pregelatinised maize starch, polyvinylpyrrolidone or hydroxypropyl methylcellulose); fillers (e.g. lactose, microcrystalline cellulose or calcium hydrogen phosphate); lubricants (e.g. magnesium stearate, talc or silica); disintegrants (e.g. potato starch or sodium starch glycolate); or wetting agents (e.g. sodium lauryl sulphate). The tablets may be coated by methods well known in the art. Liquid preparations for oral administration may take the form of, for example, solutions, syrups or suspensions, or they may be presented as a dry product for constitution with water or other suitable vehicle before use. Such liquid preparations may be prepared by conventional means with pharmaceutically acceptable additives such as suspending agents (e.g. sorbitol syrup, cellulose derivatives or hydrogenated edible fats); emulsifying agents (e.g. lecithin or acacia); non-aqueous vehicles

(e.g. almond oil, oily esters, ethyl alcohol or fractionated vegetable oils); and preservatives (e.g. methyl or propyl-p-hydroxybenzoates or sorbic acid). The preparations may also contain buffer salts, flavouring, colouring and sweetening agents as appropriate.

Preparations for oral administration may be suitably formulated to give controlled release of the active compound.

For buccal administration the composition may take the form of tablets or formulated in conventional manner.

The compounds of the invention may be formulated for parenteral administration by bolus injection or continuous infusion. Formulations for injection may be presented in unit dosage form e.g. in ampoules or in multi-dose containers, with an added preservative. The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilising and/or dispersing agents. Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle, e.g. sterile pyrogen-free water, before use.

The compounds of the invention may be formulated for topical administration in the form of ointments, creams, gels, lotions, pessaries, aerosols or drops (e.g. eye, ear or nose drops). Ointments and creams may, for example, be formulated with an aqueous or oily base with the addition of suitable thickening and/or gelling agents. Ointments for administration to the eye may be manufactured in a sterile manner using sterilised components.

Lotions may be formulated with an aqueous or oily base and will in general also contain one or more emulsifying agents, stabilising agents, dispersing agents, suspending agents, thickening agents, or colouring agents. Drops may be formulated with an aqueous or non-aqueous base also comprising one or more dispersing agents, stabilising agents, solubilising agents or suspending agents. They may also contain a preservative.

The compounds of the invention may also be formulated in rectal compositions such as suppositories or retention enemas, e.g. containing conventional suppository bases such as cocoa butter or other glycerides.

The compounds of the invention may also be formulated as depot preparations. Such long acting formulations may be administered by implantation (for example subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the compounds of the invention may be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

For intranasal administration, the compounds of the invention may be formulated as solutions for administration via a suitable metered or unitary dose device or alternatively as a powder mix with a suitable carrier for administration using a suitable delivery device.

A proposed dose of the compounds of the invention is 1 to about 1000mg per day. It will be appreciated that it may be necessary to make routine variations to the dosage, depending on the age and condition of the patient and the precise dosage will be ultimately at the discretion of the attendant physician or veterinarian. The dosage will also depend on the route of administration and the particular compound selected.

Thus for parenteral administration a daily dose will typically be in the range of 1 to about 100 mg, preferably 1 to 80 mg per day. For oral administration a daily dose will typically be within the range 1 to 300 mg e.g. 1 to 100 mg.

### EXAMPLES

In the Intermediates and Examples unless otherwise stated:

Melting points (m.p.) were determined on a Gallenkamp m.p. apparatus and are uncorrected. All temperatures refers to °C. Infrared spectra were measured on a FT-IR instrument. Proton Magnetic Resonance (<sup>1</sup>H-NMR) spectra were recorded at 400 MHz, chemical shifts are reported in ppm downfield (δ) from Me<sub>4</sub>Si, used as internal standard, and are assigned as singlets (s), doublets (d), doublets of doublets (dd), triplets (t), quartets (q) or multiplets (m). Column chromatography was carried out over silica gel (Merck AG Darmstadt, Germany). The following abbreviations are used in text: EtOAc = ethyl acetate, cHex = cyclohexane, CH<sub>2</sub>Cl<sub>2</sub> = dichloromethane, Et<sub>2</sub>O = diethyl ether, DMF = N,N'-dimethylformamide, NMP = N-methyl pyrrolidinone, DIPEA = N,N-diisopropylethylamine, DME = ethylene glycol dimethyl ether, MeOH = methanol, Et<sub>3</sub>N = triethylamine, TFA = trifluoroacetic acid, THF = tetrahydrofuran, DIBAL-H = diisobutylaluminium hydride, DMAP = dimethylaminopyridine, LHMDs = lithium hexamethyldisilazane, DMA = dimethylacetamide, NMM = N-methylmorpholine, DAST = (diethylamino)sulfur trifluoride, Tlc refers to thin layer chromatography on silica plates, and dried refers to a solution dried over anhydrous sodium sulphate; r.t. (RT) refers to room temperature.

#### Intermediate 1

##### Ethyl 2,4-dichloro-6-methyl-3-pyridinecarboxylate

The title compound was prepared according to an already published procedure: Mittelbach, Martin; *Synthesis*, 1988, 6, p.479-80.

#### Intermediate 2

##### Ethyl 2-chloro-6-methyl-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-3-pyridinecarboxylate

To a solution of 2-(1H-pyrazol-3-yl)-1,3-thiazole (7.71 g, 1.05 eq) in anh. DMF (61 mL), at 0°C, under N<sub>2</sub>, was added NaH 60% in mineral oil (2.03 g, 1.05 eq) and the reaction mixture was stirred for 10 min. at 0°C and then for 1 hr at room temperature. Intermediate 1 (11.34 g, 48.0 mmol) was then added as a solution in anh. DMF (35 mL) at 0°C and the resulting solution was heated at 110°C for 3 hr. The reaction was then quenched with water, extracted with EtOAc, washed with brine, dried over anh. Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated *in vacuo*. The crude product was purified by flash chromatography (silica gel, cHex/EtOAc 7:3) to give 7.02 g of the title compound as a white solid.

NMR (<sup>1</sup>H, CDCl<sub>3</sub>): δ 7.91 (d, 1H), 7.91 (d, 1H), 7.41 (d, 1H), 7.31 (s, 1H), 7.18 (d, 1H), 4.50 (q, 2H), 2.78 (s, 3H), 1.25 (t, 3H).

MS (m/z): 349 [MH]<sup>+</sup>.

#### Intermediate 3

##### 2-Chloro-6-methyl-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-3-pyridinylmethanol

To a solution of intermediate 2 (1.5 g, 4.3 mmol) in anh. CH<sub>2</sub>Cl<sub>2</sub> (30 mL), at -78°C, under N<sub>2</sub>, was added DIBAL-H 1.0 M in cyclohexane (12.9 mL, 3.0 eq). The reaction mixture was stirred for 1 hr at -78°C and then for 1 hr at room temperature. The reaction was then quenched with a saturated solution of Rochelle's salt, extracted with EtOAc, washed with brine, dried over anh. Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated *in vacuo*. The crude product was purified by flash chromatography (silica gel, cHex/EtOAc 1:1) to give 1.02 g of the title compound as a white solid.

NMR (<sup>1</sup>H, CDCl<sub>3</sub>): δ 8.05 (d, 1H), 7.90 (d, 1H), 7.40 (d, 1H), 7.25 (s, 1H), 7.10 (d, 1H), 4.65 (s, 2H), 4.0 (bs, 1H), 2.60 (s, 3H).

MS (m/z): 307 [M]<sup>+</sup>.

#### Intermediate 4

##### 2-chloro-6-methyl-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-3-pyridinecarbaldehyde

To a solution of intermediate 3 (150 mg, 0.5 mmol) in anh. CH<sub>2</sub>Cl<sub>2</sub> (5 mL), at room temperature, under N<sub>2</sub>, was added the Dess Martin periodinane (237 mg, 1.12 eq) and the reaction mixture was stirred for 1 hr at room temperature. The reaction was then quenched with a solution of 0.5 g of sodium thiosulfate dissolved in a saturated solution of sodium bicarbonate, extracted with EtOAc, washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated *in vacuo*. The crude product was purified by flash chromatography (silica gel, cHex/EtOAc 1:1) to give 124 mg of the title compound as a white solid.

NMR (<sup>1</sup>H, CDCl<sub>3</sub>): δ 10.4 (s, 1H), 8.0-7.9 (2d, 2H), 7.40 (2d, 2H), 7.10 (s, 1H), 2.70 (s, 3H).

MS (m/z): 305 [MH]<sup>+</sup>.

#### Intermediate 5

##### 2-Chloro-6-methyl-3-[(E)-2-(methoxy)ethenyl]-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]pyridine

To a solution of (methoxymethyl)-triphenylphosphonium chloride (4.24 g, 3 eq) in anh. THF (20 mL), at 0 °C, under N<sub>2</sub>, was added *n*-BuLi 1.6 M in cyclohexane (7.73 ml, 12.37 mmol) and the reaction mixture was brought to room temperature and then stirred for 15 min. A solution of intermediate 4 (1.25 g, 4.1 mmol) in anh. THF (15 mL) was then added and the reaction was stirred at room temperature for 1.5 hr. The reaction was then quenched with water, extracted with EtOAc, washed with brine, dried over anh. Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated *in vacuo*. The crude product was purified by flash chromatography (silica gel, cHex/EtOAc 4:1) to give 961 mg of the title compound as a white solid (E:Z = 3:2 mixture, used as such in the next step).

NMR (<sup>1</sup>H, CDCl<sub>3</sub>) principal E product : δ 7.90 (m, 1H), 7.83 (m, 1H), 7.38 (m, 1H), 7.05 (m, 1H), 7.00 (m, 1H), 6.51 (d, 1H), 5.63 (d, 1H), 3.64 (s, 3H), 2.60 (s, 3H).

MS (*m/z*): 333 [MH]<sup>+</sup>.

#### Intermediate 6

##### {2-chloro-6-methyl-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-3-pyridinyl}acetaldehyde

To a solution of intermediate 5 (938 mg, 2.8 mmol) in anh. THF (15 mL) was added 6N HCl (21 ml, 45 eq) and the reaction mixture was stirred at room temperature for 15 hr. The reaction was then quenched with sat. aq. NaHCO<sub>3</sub> until pH=7, extracted with EtOAc, washed with brine, dried over anh. Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated *in vacuo* to give 893 mg of the title compound as a white solid, which was used in the next step without further purification.

NMR (<sup>1</sup>H, CDCl<sub>3</sub>): δ 9.80 (s, 1H), 7.90-7.80 (2d, 2H), 7.70 (d, 1H), 7.20 (d, 1H), 7.0 (s, 1H), 4.25 (s, 2H), 2.70 (s, 3H).

MS (*m/z*): 319 [MH]<sup>+</sup>.

#### Intermediate 7

##### 2-{2-chloro-6-methyl-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-3-pyridinyl}ethanol

To a solution of intermediate 6 (903 mg, 2.84 mmol) in anh. MeOH (10 mL) were added CeCl<sub>3</sub> (700 mg, 1 eq) and NaBH<sub>4</sub> (107 mg, 1 eq) and the reaction mixture was stirred at room temperature for 5 min. The reaction was then quenched with water, extracted with ethyl acetate, washed with brine, dried over anh. Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated *in vacuo* to give 848 mg of the title compound as a white solid, which was used in the next step without further purification.

NMR (<sup>1</sup>H, CDCl<sub>3</sub>): δ 8.00 (m, 2H), 7.50 (d, 1H), 7.20 (m, 2H), 4.25 (t, 2H), 3.20 (t, 2H), 2.70 (s, 3H).

MS (*m/z*): 321 [MH]<sup>+</sup>.

#### Intermediate 8

##### 2-chloro-3-(2-(((1,1-dimethylethyl)(dimethyl)silyl)oxy)ethyl)-6-methyl-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]pyridine

To a solution of intermediate 7 (840 mg, 2.6 mmol) in anh. CH<sub>2</sub>Cl<sub>2</sub> (10 mL) were added 2,6-lutidine (0.67 ml, 2.2 eq) and *tert*-butyldimethylsilyl triflate (0.89 ml, 1.5 eq) and the reaction

mixture was stirred at room temperature for 15 hr. The reaction was then quenched with an aqueous solution of saturated  $\text{NH}_4\text{Cl}$ , extracted with EtOAc, washed with brine, dried over anhydrous  $\text{Na}_2\text{SO}_4$ , filtered and concentrated *in vacuo*. The product was purified by flash chromatography (silica gel,  $\text{CH}_2\text{Cl}_2/\text{EtOAc}$  3:2) to give 950 mg of the title compound as a colorless oil.

NMR ( $^1\text{H}$ ,  $\text{CDCl}_3$ ):  $\delta$  8.20 (d, 1H), 7.75 (d, 1H), 7.35 (d, 1H), 7.00 (m, 2H), 4.00 (t, 2H), 3.05 (t, 2H), 2.55 (s, 3H), 0.80 (s, 9H), -0.10 (s, 6H).

MS ( $m/z$ ): 435 [ $\text{MH}$ ] $^+$ .

#### Intermediate 9

3-(2-((1,1-dimethylethyl)(dimethyl)silyloxy)ethyl)-6-methyl-N-[6-(methoxy)-2-(trifluoromethyl)-3-pyridinyl]-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-2-pyridinamine

To a solution of Intermediate 8 (68 mg, 0.16 mmol) in anhydrous DME (2 mL) were added  $\text{Pd}_2(\text{dba})_3$  (15 mg, 0.1 eq), 2-(dicyclohexylphosphino)-2'-methylbiphenyl (17 mg, 0.3 eq),  $\text{K}_3\text{PO}_4$  (90 mg, 3 eq) and intermediate 34 (37 mg, 1.2 eq) and the reaction mixture was submitted to microwave irradiation (150W, 100°C, 60 psi) for two 20 min cycles. The reaction was then quenched with an aqueous solution of saturated  $\text{NH}_4\text{Cl}$ , extracted with EtOAc, washed with brine, dried over anhydrous  $\text{Na}_2\text{SO}_4$ , filtered and concentrated *in vacuo*. The product was purified by flash chromatography (silica gel,  $\text{CH}_2\text{Cl}_2/\text{MeOH}$  99:1) to give 22 mg of the title compound as a yellow foam.

NMR ( $^1\text{H}$ ,  $\text{CDCl}_3$ ):  $\delta$  8.40 (d, 1H), 7.85 (bs, 1H), 7.75 (bs, 1H), 7.31 (s, 1H), 7.03 (s, 1H), 6.90 (d, 1H), 6.63 (s, 1H), 4.16 (t, 2H), 3.94 (s, 3H), 2.76 (t, 2H), 2.36 (s, 3H), 0.77 (s, 9H), 0.04 (s, 6H).

#### Intermediate 10

3-(2-((1,1-Dimethylethyl)(dimethyl)silyloxy)ethyl)-6-methyl-N-[2-methyl-6-(methoxy)-3-pyridinyl]-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-2-pyridinamine

As in intermediate 9, except that intermediate 75 (2-methyl-6-(methoxy)-3-pyridinamine) was used instead of intermediate 34 (6-(Methoxy)-2-(trifluoromethyl)-3-pyridinamine).

MS ( $m/z$ ): 537 [ $\text{MH}$ ] $^+$ .

#### Intermediate 11

N-[2,6-Bis(methoxy)-3-pyridinyl]-3-(2-((1,1-dimethylethyl)(dimethyl)silyloxy)ethyl)-6-methyl-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-2-pyridinamine

As in intermediate 9, except that 2,6-bis(methoxy)-3-pyridinamine was used instead of intermediate 34 (6-(Methoxy)-2-(trifluoromethyl)-3-pyridinamine).

NMR ( $^1\text{H}$ ,  $\text{CDCl}_3$ ):  $\delta$  8.70 (d, 1H); 7.86 (d, 1H); 7.75 (d, 1H); 7.32 (d, 1H); 7.03 (d, 1H); 6.60 (s, 1H); 6.32 (d, 1H); 4.15 (t, 2H); 3.99 (s, 3H); 3.88 (s, 3H); 2.47 (bs, 1H); 2.77 (t, 2H); 2.46 (s, 3H); 0.82 (s, 12H).

MS ( $m/z$ ): 553 [ $\text{MH}$ ] $^+$ .

Intermediate 12

*N*<sup>6</sup>-[3-(2-[(1,1-Dimethylethyl)(dimethyl)silyloxy]ethyl)-6-methyl-4-[3-(1,3-thiazol-2-yl)-1*H*-pyrazol-1-yl]-2-pyridinyl]-*N*<sup>6</sup>,*N*<sup>6</sup>,4-trimethyl-2,5-pyridinediamine

As in intermediate 9, except that intermediate 36 (*N*<sup>6</sup>,*N*<sup>6</sup>,4-trimethyl-2,5-pyridinediamine) was used instead of intermediate 34 (6-(Methyloxy)-2-(trifluoromethyl)-3-pyridinamine).

NMR (<sup>1</sup>H, DMSO-*d*<sub>6</sub>): δ 8.35 (d, 1H), 8.00 (s, 1H), 7.99 (d, 1H), 7.87 (s, 1H), 7.83 (d, 1H), 7.07 (d, 1H), 6.69 (s, 1H), 6.63 (s, 1H), 3.95 (t, 2H), 3.10 (s, 6H), 2.98 (t, 2H), 2.28 (s, 3H), 2.18 (s, 3H), 0.85 (s, 9H), 0.00 (s, 6H).

MS (*m/z*): 550 [M-H]<sup>+</sup>.

Intermediate 13

*N*-[2-(Difluoromethyl)-4-(methyloxy)phenyl]-3-(2-[(1,1-dimethylethyl)(dimethyl)silyloxy]ethyl)-6-methyl-4-[3-(1,3-thiazol-2-yl)-1*H*-pyrazol-1-yl]-2-pyridinamine

As in intermediate 9, except that intermediate 40 (2-(difluoromethyl)-4-(methyloxy)aniline) was used instead of intermediate 34 (6-(Methyloxy)-2-(trifluoromethyl)-3-pyridinamine).

NMR (<sup>1</sup>H, CDCl<sub>3</sub>): δ 7.9 (d, 1H), 7.8 (d, 1H), 7.68 (bs, 1H), 7.6 (d, 1H), 7.38 (d, 1H), 7.17 (d, 1H), 7.08 (d, 1H), 7.04 (dd, 1H), 6.77 (t, 1H; *J*<sub>H-F</sub> = 55.6 Hz), 6.64 (s, 1H), 4.24 (t, 2H), 3.89 (s, 3H), 2.81 (t, 2H), 2.41 (s, 3H), 0.88 (s, 9H), 0.00 (s, 6H).

MS (*m/z*): 572 [M-H]<sup>+</sup>.

Intermediate 14

*N*-[2-Chloro-4-(methyloxy)phenyl]-3-(2-[(1,1-dimethylethyl)(dimethyl)silyloxy]ethyl)-6-methyl-4-[3-(1,3-thiazol-2-yl)-1*H*-pyrazol-1-yl]-2-pyridinamine

As in intermediate 9, except that intermediate 41 (2-chloro-4-(methyloxy)aniline) was used instead of intermediate 34 (6-(Methyloxy)-2-(trifluoromethyl)-3-pyridinamine).

NMR (<sup>1</sup>H, CDCl<sub>3</sub>): δ 8.37 (d, 1H), 7.86 (d, 1H), 7.82 (bs, 1H), 7.77 (d, 1H), 7.32 (d, 1H), 7.03 (d, 1H), 6.95 (d, 1H), 6.84-6.82 (dd, 1H), 6.65 (s, 1H), 4.19 (t, 2H), 3.79 (s, 3H), 2.81 (t, 2H), 2.48 (s, 3H), 0.83 (s, 9H), 0.00 (s, 6H).

Intermediate 15

*N*-[2,4-Bis(methyloxy)phenyl]-3-(2-[(1,1-dimethylethyl)(dimethyl)silyloxy]ethyl)-6-methyl-4-[3-(1,3-thiazol-2-yl)-1*H*-pyrazol-1-yl]-2-pyridinamine

As in intermediate 9, except that 2,4-bis(methyloxy)aniline was used instead of intermediate 34 (6-(Methyloxy)-2-(trifluoromethyl)-3-pyridinamine).

NMR (<sup>1</sup>H, CDCl<sub>3</sub>): δ 8.21 (d, 1H), 7.84 (d, 1H), 7.75 (d, 1H), 7.34 (s, 1H), 7.31 (d, 1H), 7.02 (d, 2H), 6.50 (s, 1H), 3.96 (t, 2H), 3.78 (s, 3H), 3.72 (s, 3H), 3.05 (t, 2H), 2.55 (s, 3H), 1.81 (s, 9H), 0.00 (s, 6H).

MS (*m/z*): 553 [M-H]<sup>+</sup>.

Intermediate 16



4-((3-(2-((1,1-dimethylethyl)(dimethyl)silyloxy)ethyl)-6-methyl-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-2-pyridinyl)amino)-3-((trifluoromethyl)oxy)benzonitrile

As in intermediate 9, except that 4-amino-3-((trifluoromethyl)oxy)benzonitrile was used instead of intermediate 34 (6-(Methyloxy)-2-(trifluoromethyl)-3-pyridinamine).

MS (*m/z*): 601 [MH]<sup>+</sup>.

Intermediate 17

4-((3-(2-((1,1-Dimethylethyl)(dimethyl)silyloxy)ethyl)-6-methyl-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-2-pyridinyl)amino)-3-ethylbenzonitrile

As in intermediate 9, except that 4-amino-3-ethylbenzonitrile was used instead of intermediate 34 (6-(Methyloxy)-2-(trifluoromethyl)-3-pyridinamine).

NMR (<sup>1</sup>H, CDCl<sub>3</sub>): δ 8.37 (bs, 1H), 8.07 (d, 1H), 7.87 (d, 1H), 7.77 (d, 1H), 7.43 (bs, 1H), 7.3 (d, 1H), 7.06 (d, 1H), 6.78 (s, 1H), 4.31 (m, 2H), 2.80 (m, 2H), 2.67 (q, 2H), 2.50 (s, 3H), 2.46 (m, 1H), 1.28 (t, 3H).

MS (*m/z*): 432 [MH]<sup>+</sup>.

Intermediate 18

3-(2-((1,1-Dimethylethyl)(dimethyl)silyloxy)ethyl)-N-(4-fluoro-2-methylphenyl)-6-methyl-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-2-pyridinamine

As in intermediate 9, except that 4-fluoro-2-methylaniline was used instead of intermediate 34 (6-(Methyloxy)-2-(trifluoromethyl)-3-pyridinamine).

NMR (<sup>1</sup>H, CDCl<sub>3</sub>): δ 7.86 (s, 1H), 7.74 (d, 1H), 7.55 (d, 1H), 7.32 (s, 1H), 7.03 (s, 1H), 6.87 (m, 2H), 6.58 (s, 1H), 4.23 (t, 2H), 2.93 (bs, 1H), 2.76 (t, 2H), 2.40 (s, 3H), 2.24 (s, 3H), 0.83 (s, 9H), 0.02 (s, 6H).

MS (*m/z*): 524 [MH]<sup>+</sup>.

Intermediate 19

3-(2-((1,1-Dimethylethyl)(dimethyl)silyloxy)ethyl)-6-methyl-N-[2-methyl-4-(1H-pyrazol-1-yl)phenyl]-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-2-pyridinamine

As in intermediate 9, except that intermediate 43 (2-methyl-4-(1H-pyrazol-1-yl)aniline) was used instead of intermediate 34 (6-(Methyloxy)-2-(trifluoromethyl)-3-pyridinamine).

NMR (<sup>1</sup>H, DMSO-d<sub>6</sub>): δ 8.49 (dd, 1H), 8.36 (d, 1H), 8.03 (s, 1H), 7.99 (d, 1H), 7.84 (d, 1H), 7.77 (m, 2H), 7.70 (d, 1H), 7.65 (dd, 1H), 7.08 (d, 1H), 6.84 (s, 1H), 6.59 (m, 1H), 4.04 (t, 2H), 2.99 (t, 2H), 2.38 (s, 3H), 2.35 (s, 3H), 0.84 (s, 9H), 0.00 (s, 6H).

MS (*m/z*): 572 [MH]<sup>+</sup>.

Intermediate 20

3-(2-((1,1-Dimethylethyl)(dimethyl)silyloxy)ethyl)-N-[4-(nitro)-2-(trifluoromethyl)phenyl]-6-methyl-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-2-pyridinamine

As in intermediate 9, except that 4-nitro-2-(trifluoromethyl)aniline was used instead of intermediate 34 (6-(Methyloxy)-2-(trifluoromethyl)-3-pyridinamine).

NMR ( $^1\text{H}$ ,  $\text{CDCl}_3$ ):  $\delta$  8.37(s, 1H), 8.21(d, 1H), 8.18(d, 1H), 8.15(d, 1H), 7.87(d, 1H), 7.34(d, 1H), 7.03(d, 1H), 6.75(s, 1H), 6.72(s, 1H), 3.98(t, 2H), 3.08(t, 2H), 2.55(s, 3H), 1.55(s, 6H), 0.75(s, 9H).  
MS ( $m/z$ ): 605  $[\text{MH}]^+$ .

#### Intermediate 21

2-[6-Methyl-2-[[6-(methyloxy)-2-(trifluoromethyl)-3-pyridinyl]amino]-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-3-pyridinyl]ethanol

To a solution of intermediate 9 (240 mg, 0.38 mmol) in anhyd. THF (5 mL) was added  $\text{Et}_3\text{N} \cdot 3\text{HF}$  (0.187 mL, 3 eq) and the reaction mixture was stirred for 15 hr at r.t.. The reaction was then quenched with sat. aq.  $\text{NH}_4\text{Cl}$ , extracted with EtOAc, washed with sat. aq. NaCl, dried over anhyd.  $\text{Na}_2\text{SO}_4$ , filtered and concentrated *in vacuo*. The product was purified by flash chromatography (silica gel,  $\text{cHex}/\text{EtOAc}$  1:1) to give 180 mg of the title compound as a colorless oil.

NMR ( $^1\text{H}$ ,  $\text{CDCl}_3$ ):  $\delta$  8.45 (bs, 1H), 8.20 (d, 1H), 7.85 (d, 1H), 7.85 (2d, 2H), 7.65 (dd, 1H), 7.30 (d, 1H), 7.05 (d, 1H), 6.85 (s, 1H), 4.20 (t, 2H), 2.85 (t, 2H), 2.50 (s, 3H).  
MS ( $m/z$ ): 514  $[\text{MH}]^+$ .

#### Intermediate 22

2-[6-Methyl-2-[[6-methyl-2-(methyloxy)-3-pyridinyl]amino]-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-3-pyridinyl]ethanol

As in intermediate 21, except that intermediate 10 was used instead of intermediate 9.

MS ( $m/z$ ): 423  $[\text{MH}]^+$ .

#### Intermediate 23

2-[2-[[6-Bis(methyloxy)-3-pyridinyl]amino]-6-methyl-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-3-pyridinyl]ethanol

As in intermediate 21, except that intermediate 11 was used instead of intermediate 9.

NMR ( $^1\text{H}$ ,  $\text{CDCl}_3$ ):  $\delta$  8.60 (d, 1H); 7.88 (d, 1H); 7.74 (d, 1H); 7.31 (d, 1H); 7.02 (d, 1H); 6.56 (s, 1H); 6.32 (d, 1H); 4.13 (t, 2H); 3.97 (s, 3H); 3.86 (s, 3H); 3.78 (t, 1H); 2.77 (t, 2H); 2.46 (s, 3H).

MS ( $m/z$ ): 439  $[\text{MH}]^+$ .

#### Intermediate 24

2-[2-[[6-(Dimethylamino)-4-methyl-3-pyridinyl]amino]-6-methyl-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-3-pyridinyl]ethanol

As in intermediate 21, except that intermediate 12 was used instead of intermediate 9.

NMR ( $^1\text{H}$ ,  $\text{DMSO}-d_6$ ):  $\delta$  8.26 (d, 1H), 8.02 (s, 1H), 7.97 (s, 1H), 7.90 (d, 1H), 7.73 (d, 1H), 6.98 (d, 1H), 6.62 (s, 1H), 6.53 (s, 1H), 5.16 (t, 1H), 3.81 (q, 2H), 3.00 (s, 6H), 2.21 (s, 3H), 2.10 (s, 3H).

MS ( $m/z$ ): 436  $[\text{MH}]^+$ .

Intermediate 25

2-{2-([2-(Difluoromethyl)-4-(methoxy)phenyl]amino)-6-methyl-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-3-pyridinyl}ethanol

As in intermediate 21, except that intermediate 13 was used instead of intermediate 9.

NMR (<sup>1</sup>H, CDCl<sub>3</sub>): δ 7.88 (d, 1H), 7.76 (d, 1H), 7.67 (d, 1H), 7.48 (bs, 1H), 7.34 (d, 1H), 7.09 (d, 1H), 7.06 (d, 1H), 7.01 (dd, 1H), 6.74 (t, 1H; J<sub>H-F</sub> = 55.3 Hz), 6.62 (s, 1H), 4.21 (m, 2H), 3.85 (s, 3H), 3.18 (t, 1H), 2.83 (t, 2H), 2.4 (s, 3H).

MS (m/z): 458 [MH]<sup>+</sup>.

Intermediate 26

2-{2-([2-Chloro-4-(methoxy)phenyl]amino)-6-methyl-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-3-pyridinyl}ethanol

As in intermediate 21, except that intermediate 14 was used instead of intermediate 9.

NMR (<sup>1</sup>H, CDCl<sub>3</sub>): δ 8.27 (d, 1H), 7.86 (d, 1H), 7.58 (bs, 1H), 7.33 (d, 1H), 7.57 (d, 1H), 7.05 (d, 1H), 6.95 (d, 1H), 6.84-6.82 (dd, 1H), 6.64 (s, 1H), 4.20 (t, 2H), 3.78 (s, 3H), 3.49 (t, 1H), 2.8 (t, 2H), 2.48 (s, 3H).

Intermediate 27

2-{2-([2,4-Bis(methoxy)phenyl]amino)-6-methyl-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-3-pyridinyl}ethanol

As in intermediate 21, except that intermediate 15 was used instead of intermediate 9.

NMR (<sup>1</sup>H, CDCl<sub>3</sub>): δ 8.35 (d, 1H), 7.85 (d, 1H), 7.73 (d, 1H), 7.40 (s, 1H), 7.31 (d, 1H), 7.05 (d, 1H), 6.52 (s, 1H), 6.50 (s, 1H), 4.18 (m, 2H), 3.87 (s, 3H), 3.80 (s, 3H), 2.85 (t, 2H), 2.48 (s, 3H).

MS (m/z): 440 [MH]<sup>+</sup>.

Intermediate 28

4-([3-(2-hydroxyethyl)-6-methyl-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-2-pyridinyl]amino)-3-[(trifluoromethyl)oxy]benzonitrile

As in intermediate 21, except that intermediate 16 was used instead of intermediate 9.

NMR (<sup>1</sup>H, CDCl<sub>3</sub>): δ 9.27 (s, 1H); 8.45 (d, 1H); 7.90 (d, 1H); 7.80 (d, 1H); 7.50 (dd, 2H); 7.30 (d, 1H); 7.10 (d, 1H); 6.95 (d, 1H); 4.45 (t, 2H); 2.80 (t, 2H); 2.50 (s, 3H); 2.45 (s, 1H).

MS (m/z): 487 [MH]<sup>+</sup>.

Intermediate 29

3-ethyl-4-([3-(2-hydroxyethyl)-6-methyl-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-2-pyridinyl]amino)benzonitrile

As in intermediate 21, except that intermediate 17 was used instead of intermediate 9.

NMR ( $^1\text{H}$ ,  $\text{CDCl}_3$ ):  $\delta$  8.37 (bs, 1H), 8.07 (d, 1H), 7.87 (d, 1H), 7.77 (d, 1H), 7.43 (bs, 1H), 7.3 (d, 1H), 7.06 (d, 1H), 6.78 (s, 1H), 4.31 (m, 2H), 2.80 (m, 2H), 2.67 (q, 2H), 2.50 (s, 3H), 2.46 (m, 1H), 1.28 (t, 3H).

MS ( $m/z$ ): 414  $[\text{MH}]^+$ .

#### Intermediate 30

2-[2-[(4-fluoro-2-methylphenyl)amino]-6-methyl-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-3-pyridinyl]ethanol

As in intermediate 21, except that intermediate 18 was used instead of intermediate 9.

NMR ( $^1\text{H}$ ,  $\text{CDCl}_3$ ):  $\delta$  7.92 (d, 1H), 7.70 (s, 1H), 7.65 (d, 1H), 7.50 (s, 1H), 7.32 (s, 1H), 7.00 (s, 1H), 6.80-6.95 (m, 2H), 6.58 (s, 1H), 4.25 (bs, 2H), 2.93 (bs, 1H), 2.83 (t, 2H), 2.40 (s, 3H), 2.27 (s, 3H).

MS ( $m/z$ ): 410  $[\text{MH}]^+$ .

#### Intermediate 31

2-[6-Methyl-2-[(2-methyl-4-(1H-pyrazol-1-yl)phenyl)amino]-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-3-pyridinyl]ethanol

As in intermediate 21, except that intermediate 19 was used instead of intermediate 9.

NMR ( $^1\text{H}$ ,  $\text{DMSO}-d_6$ ):  $\delta$  8.58 (s, 1H), 8.38 (dd, 1H), 8.28 (d, 1H), 7.90 (d, 1H), 7.74 (m, 1H), 7.68 (dd, 1H), 7.66 (d, 1H), 7.57 (dd, 1H), 7.00 (d, 1H), 6.81 (s, 1H), 6.49 (m, 1H), 5.61 (bs, 1H), 3.94 (m, 2H), 2.70 (m, 2H), 2.33 (s, 3H), 2.28 (s, 3H).

MS ( $m/z$ ): 458  $[\text{MH}]^+$ .

#### Intermediate 32

2-[2-[(4-nitro-2-(trifluoromethyl)phenyl)amino]-6-methyl-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-3-pyridinyl]ethanol

As in intermediate 21, except that intermediate 20 was used instead of intermediate 9.

NMR ( $^1\text{H}$ ,  $\text{CDCl}_3$ ):  $\delta$  9.05 (s, 1H), 8.50 (d, 1H), 8.25 (d, 1H), 8.15 (dd, 1H), 7.85 (d, 1H), 7.75 (d, 1H), 7.32 (d, 1H), 7.05 (d, 1H), 6.90 (s, 1H), 4.20 (m, 2H), 2.80 (t, 2H), 2.60 (bs, 1H), 2.55 (s, 3H).

MS ( $m/z$ ): 491  $[\text{MH}]^+$ .

#### Intermediate 33

6-(methyloxy)-3-nitro-2-(trifluoromethyl)pyridine

To a solution of 2-chloro-3-nitro-5-methoxypyridine (500 mg, 2.6 mmol) in DMA (5 mL), at r.t., under  $\text{N}_2$ , were added Cu (1 g, 6 eq) and  $\text{CF}_3\text{Br}_2$  (500  $\mu\text{L}$ , large excess). The reaction mixture was stirred at  $100^\circ\text{C}$  for 8 hr. The brown slurry obtained was extracted with  $\text{Et}_2\text{O}$  (3x50 mL). The combined organic extracts were dried under vacuum and the crude product was purified by flash chromatography (silica gel,  $\text{CH}_2\text{Cl}_2$  100%) to give the title compound (316 mg, 53%) as a brown oil.

NMR ( $^1\text{H}$ ,  $\text{CDCl}_3$ ):  $\delta$  8.35 (d, 1H), 7.2 (d, 1H), 4.3 (s, 3H).

Intermediate 346-(Methoxy)-2-(trifluoromethyl)-3-pyridinamine

To a solution of intermediate 33 (98 mg, 0.44 mmol) in MeOH (10 mL), at r.t., was added Pd/C 10% (30 mg 30% w/w) and the resulting suspension subjected to hydrogenation (1 atm) for 2 hr. The palladium was filtered off and the solvent was evaporated under reduced pressure. The crude product was purified by flash chromatography (silica gel, CH<sub>2</sub>Cl<sub>2</sub>/MeOH 95:5) to give the title compound (37 mg, 44%) as a red oil.

NMR (<sup>1</sup>H, CDCl<sub>3</sub>): δ 7.10 (d, 1H); 6.56 (d, 1H); 3.85 (s, 3H); 3.82-3.55 (bs, 2H).

Intermediate 35N,N,4-Trimethyl-5-nitro-2-pyridinamine

To 2-chloro-4-methyl-5-nitropyridine (2.78 g, 16.1 mmol), at r.t., under N<sub>2</sub>, was added Me<sub>2</sub>NH 2N/THF (55.4 mL). The reaction mixture was heated at 80°C for 2 hr. It was cooled down to r.t. and partitioned between CH<sub>2</sub>Cl<sub>2</sub> and water. The phases were separated and the aqueous layer was further extracted with CH<sub>2</sub>Cl<sub>2</sub> (3x30 mL). The combined organic extracts were dried over anh. Na<sub>2</sub>SO<sub>4</sub>, the solids were filtered and the solvent evaporated. The crude title compound was used in the next step without further purification (2.97 g, quantitative yield).

NMR (<sup>1</sup>H, CDCl<sub>3</sub>): δ 8.99 (s, 1H), 6.24 (s, 1H), 3.2 (s, 6H), 2.62 (s, 3H).

MS (m/z): 182 [MH]<sup>+</sup>.

Intermediate 36N<sup>2</sup>,N<sup>2</sup>,4-Trimethyl-2,5-pyridinediamine

To a suspension of intermediate 35 (2.97 g, 16.31 mmol) in a 1:1 mixture of MeOH/H<sub>2</sub>O (100 mL), at r.t., under N<sub>2</sub>, were added Fe (3.19 g, 3.5 eq) and NH<sub>4</sub>Cl (3.04 g, 3.5 eq). The reaction mixture was heated at 80°C for 1.5 hr. The mixture was filtered and the solid was washed with MeOH. The crude was evaporated to dryness and partitioned between CH<sub>2</sub>Cl<sub>2</sub> and water. The phases were separated and the aqueous layer was further extracted with CH<sub>2</sub>Cl<sub>2</sub> (3x30 mL). The combined organic extracts were dried over anh. Na<sub>2</sub>SO<sub>4</sub>, the solids were filtered and the solvent evaporated. The crude product was purified by flash-chromatography (silica gel, EtOAc/MeOH 9:1 → 7:3) to give the title compound as a yellow solid (340 mg, 14%).

NMR (<sup>1</sup>H, DMSO-d<sub>6</sub>): δ 7.52 (s, 1H), 6.36 (s, 1H), 4.17 (bs, 2H), 2.83 (s, 6H), 2.03 (s, 3H).

MS (m/z): 152 [MH]<sup>+</sup>.

Intermediate 37(5-Methoxy-2-nitro-phenyl)-methanol

To a suspension of cyanuric chloride (1.84 g, 1 eq) in anh. DME (60 mL) at r.t., under N<sub>2</sub>, was added NMM (1.1 mL, 1 eq). The reaction mixture was stirred for 2 min and a precipitate was formed. A solution of 5-(methoxy)-2-nitrobenzoic acid (2.0 g, 10 mmol) in anh. DME (20 mL) was added and the reaction mixture was stirred for 4 hr. The suspension was filtered

and a solution of  $\text{NaBH}_4$  (0.57 g, 1.5 eq) in water (30 mL) was added at  $0^\circ\text{C}$ . The reaction mixture was stirred for 20 min at  $0^\circ\text{C}$ . It was then diluted with  $\text{Et}_2\text{O}$  (10 mL) and acidified to  $\text{pH}=5$  by addition of sat. aq.  $\text{NH}_4\text{Cl}$ . The phases were separated and the aqueous layer was extracted with  $\text{Et}_2\text{O}$  (2x 100mL). The combined organic extracts were washed with sat. aq.  $\text{Na}_2\text{CO}_3$  and dried over anh.  $\text{Na}_2\text{SO}_4$ . The solids were filtered and the solvent evaporated to dryness. The crude product was purified by flash chromatography (silica gel,  $\text{cHex/EtOAc}$  8:2) to give the title compound (958 mg, 53 %).

NMR ( $^1\text{H}$ ,  $\text{CDCl}_3$ ):  $\delta$  8.15 (d, 1H), 7.19 (m, 1H), 6.85 (dd, 1H), 4.95 (d, 2H), 3.89 (s, 3H), 2.5 (t, 1H).

#### Intermediate 38

##### 5-(Methoxy)-2-nitrobenzaldehyde

To a solution of intermediate 37 (1.44 g, 7.9 mmol) in anh.  $\text{CH}_2\text{Cl}_2$  (40 mL) at r.t., under  $\text{N}_2$ , was added Dess-Martin periodinane (3.68 g, 1.1 eq). The reaction mixture was stirred for 3 hr at r.t., then sat. aq.  $\text{Na}_2\text{S}_2\text{O}_3$  (5 mL) and sat. aq.  $\text{NaHCO}_3$  (20 mL) were added. The phases were separated and the aqueous layer was extracted with  $\text{EtOAc}$  (2x100 mL). The combined organic extracts were dried over anh.  $\text{Na}_2\text{SO}_4$ , the solids were filtered and the solvent evaporated to dryness to give 1.45 g (100%) of the title compound.

NMR ( $^1\text{H}$ ,  $\text{CDCl}_3$ ):  $\delta$  10.47 (s, 1H), 8.14 (d, 1H), 7.31 (d, 1H), 7.13 (dd, 1H), 3.94 (s, 3H).

#### Intermediate 39

##### 2-(Difluoromethyl)-4-(methoxy)-1-nitrobenzene

To a solution of intermediate 38 (250 mg, 1.38 mmol) in anh.  $\text{CH}_2\text{Cl}_2$  (10 mL), at  $-78^\circ\text{C}$ , under  $\text{N}_2$ , was added slowly DAST (2x0.4 mL, 2.2 eq). The reaction mixture was stirred at r.t. for 1.5 hr, after which was added sat. aq.  $\text{NaCl}$ . The phases were separated and the aqueous layer was extracted with  $\text{EtOAc}$  (3x20 mL). The combined organic extracts were dried over anh.  $\text{Na}_2\text{SO}_4$ , the solids were filtered and the solvent was evaporated to dryness. The crude product was purified by flash chromatography (silica gel,  $\text{cHex/EtOAc}$  8:2) to give the title compound (176 mg, 63 %) as a yellow oil.

NMR ( $^1\text{H}$ ,  $\text{CDCl}_3$ ):  $\delta$  8.21 (d, 1H), 7.43 (t, 1H;  $J_{\text{H-F}} = 54.9$  Hz), 7.33 (d, 1H), 7.06 (dd, 1H), 3.95 (s, 3H).

#### Intermediate 40

##### 2-(Difluoromethyl)-4-(methoxy)aniline

To a solution of intermediate 39 (176 mg, 0.87 mmol) in anh.  $\text{MeOH}$  (8.7 mL), at r.t., under  $\text{N}_2$ , was added  $\text{Pd/C}$  10% (88 mg, 5% wt). The reaction mixture was placed under an atmosphere of  $\text{H}_2$  for 5 hr. The catalyst was filtered off and the solution obtained was evaporated to dryness. The crude product was purified by flash chromatography (silica gel,  $\text{cHex/EtOAc}$  9:1) to give the title compound (27 mg, 20 %) as a yellow oil.

NMR ( $^1\text{H}$ ,  $\text{CDCl}_3$ ):  $\delta$  6.88-6.80 (m, 2H); 6.7-6.67 (m, 1H); 6.62 (t, 1H;  $J_{\text{H-F}} = 55.6$  Hz); 3.8-3.5 (bs, 2H); 3.76 (s, 3H)

MS (*m/z*): 174 [MH]<sup>+</sup>.

#### Intermediate 41

##### 2-Chloro-4-(methoxy)aniline

To a suspension of 4-amino-3-chlorophenol hydrochloride (500 mg, 2.7 mmol) in acetone (5 mL), at r.t., under N<sub>2</sub>, were added K<sub>2</sub>CO<sub>3</sub> (138 mg, 1 eq) and *t*-BuOK (311 mg, 1 eq). MeI was then added (173  $\mu$ L, 1 eq.). The reaction mixture was stirred at r.t. for 2 hr. CH<sub>2</sub>Cl<sub>2</sub> (10 mL) was added and the reaction quenched with H<sub>2</sub>O (5 mL). The phases were separated and the aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub> (5 mL). The combined organic extracts were dried over anhyd. Na<sub>2</sub>SO<sub>4</sub>, the solids were filtered and the solvent evaporated under reduced pressure. The crude product was purified by flash chromatography (silica gel, CH<sub>2</sub>Cl<sub>2</sub> 100%) to give the title compound (180 mg, 41 %).

NMR (<sup>1</sup>H, DMSO):  $\delta$  6.81 (d, 1H), 6.72 (s, 1H), 6.68 (d, 1H), 4.81 (s, 2H), 3.63 (s, 3H).

#### Intermediate 42

##### 1,1-Dimethylethyl (4-bromo-2-methylphenyl)carbamate

To a solution of 4-bromo-2-methylaniline (1 g, 5.37 mmol) in 1,4-dioxane (11 mL) and H<sub>2</sub>O (4 mL), at r.t., were added Et<sub>3</sub>N (2.7 mL, 1.2 eq) and BOC<sub>2</sub>O (4.2 g, 1.2 eq). The reaction mixture was stirred at r.t. for 96 hr. Saturated aq. NH<sub>4</sub>Cl and EtOAc (20 mL) were added and the phases were separated. The aqueous layer was further extracted with EtOAc (2x20 mL). The combined organic extracts were dried over anhyd. Na<sub>2</sub>SO<sub>4</sub>, the solids were filtered and the solvent evaporated. The residue was purified by SCX Column (Eluents: CH<sub>2</sub>Cl<sub>2</sub>, MeOH and NH<sub>3</sub> (0.5 M in MeOH)) to give the title compound as a white solid (1.22 g, 79%).

NMR (<sup>1</sup>H, DMSO-*d*<sub>6</sub>):  $\delta$  8.55 (s, 1H), 7.35 (m, 1H), 7.28 (m, 2H), 2.17 (s, 3H), 1.44 (s, 9H).

MS (*m/z*): 230 [MH-*t*Bu]<sup>+</sup>; 186 [MH-BOC]<sup>+</sup>.

#### Intermediate 43

##### 2-Methyl-4-(1H-pyrazol-1-yl)aniline

A solution of intermediate 42 (200 mg, 0.7 mmol), 1H-pyrazole (95 mg, 2 eq), CuI (133 mg, 1 eq), K<sub>2</sub>CO<sub>3</sub> (290 mg, 2.1 eq) and (1R,2R)-diaminomethylcyclohexane (100 mg, 1 eq) in anhyd. NMP (1 mL), under N<sub>2</sub>, was heated at 150°C for 6 h. It was cooled down to r.t. and poured into water. EtOAc was added and the phases were separated. The aqueous layer was further extracted with EtOAc (2x10 mL). The combined organic extracts were dried over anhyd. Na<sub>2</sub>SO<sub>4</sub>, the solids were filtered and the solvent evaporated. The residue was purified by flash chromatography (silica gel, *n*-Hex/EtOAc 8:2) to give the title compound as a white solid (85.6 mg, 70%).

NMR (<sup>1</sup>H, CDCl<sub>3</sub>):  $\delta$  7.77 (dd, 1H), 7.66 (d, 1H), 7.39 (d, 1H), 7.29 (dd, 1H), 6.72 (d, 1H), 6.40 (t, 1H), 2.85 (bs, 1H).

MS (*m/z*): 174 [MH]<sup>+</sup>.

#### Intermediate 44

Ethyl 4-(1'-H-1,3'-bipyrazol-1'-yl)-2-chloro-6-methyl-3-pyridinecarboxylate

As in intermediate 2, except that intermediate 58 was used instead of 2-(1'-H-pyrazol-3-yl)-1,3-thiazole.

NMR (<sup>1</sup>H, CDCl<sub>3</sub>): δ 8.10 (s, 1H), 7.90 (d, 1H), 7.68 (s, 1H), 7.29 (s, 1H), 6.85 (d, 1H), 6.43 (s, 1H), 4.40 (t, 2H), 2.57 (s, 3H), 1.24 (q, 3H).

MS (m/z): 332 [MH]<sup>+</sup>.

Intermediate 45[4-(1'-H-1,3'-Bipyrazol-1'-yl)-2-chloro-6-methyl-3-pyridinyl]methanol

As in intermediate 3, except that intermediate 44 was used instead of intermediate 2.

NMR (<sup>1</sup>H, CDCl<sub>3</sub>): δ 7.12 (s, 1H), 7.99 (d, 1H), 7.74 (s, 1H), 7.23 (s, 1H), 6.84 (d, 1H), 6.44 (s, 1H), 4.76 (bs, 2H), 3.74 (bs, 1H), 2.60 (s, 3H).

MS (m/z): 290 [MH]<sup>+</sup>.

Intermediate 464-(1'-H-1,3'-Bipyrazol-1'-yl)-2-chloro-6-methyl-3-pyridinecarbaldehyde

As in intermediate 4, except that intermediate 45 was used instead of intermediate 3.

NMR (<sup>1</sup>H, CDCl<sub>3</sub>): δ 10.45 (s, 1H), 8.13 (s, 1H), 7.94 (s, 1H), 7.71 (s, 1H), 7.33 (s, 1H), 6.82 (s, 1H), 6.41 (s, 1H), 2.67 (s, 3H).

MS (m/z): 288 [MH]<sup>+</sup>.

Intermediate 471'-{2-Chloro-6-methyl-3-[(E)-2-(methyloxy)ethenyl]-4-pyridinyl}-1'-H-1,3'-bipyrazole

As in intermediate 5, except that intermediate 46 was used instead of intermediate 4.

NMR (<sup>1</sup>H, CDCl<sub>3</sub>): δ 8.1-7.8 (m, 2H), 7.7-7.6 (m, 2H), 7.5-7.2 (m, 1H), 7.0-6.8 (s, 1H), 6.1 (d, 1H), 5.4 (s, 1H), 3.40 (s, 3H), 2.60 (s, 3H).

MS (m/z): 316 [MH]<sup>+</sup>.

Intermediate 48[4-(1'-H-1,3'-Bipyrazol-1'-yl)-2-chloro-6-methyl-3-pyridinyl]acetaldehyde

As in intermediate 6, except that intermediate 47 was used instead of intermediate 5.

MS (m/z): 302 [MH]<sup>+</sup>.

Intermediate 492-[4-(1'-H-1,3'-Bipyrazol-1'-yl)-2-chloro-6-methyl-3-pyridinyl]ethanol

As in intermediate 7, except that intermediate 48 was used instead of intermediate 6.

NMR (<sup>1</sup>H, CDCl<sub>3</sub>): δ 7.9 (d, 1H), 8.0 (d, 1H), 7.4 (t, 1H), 7.1 (s, 1H), 6.8 (d, 2H), 4.2 (t, 2H), 3.90 (tb, 1H), 3.2 (t, 2H), 2.40 (s, 3H).

MS (m/z): 304 [MH]<sup>+</sup>.



Intermediate 50

1'-[2-Chloro-3-(2-[(1,1-dimethylethyl)(dimethyl)silyloxy]ethyl)-6-methyl-4-pyridinyl]-1''H-1,3'-bipyrazole

As in Intermediate 8, except that intermediate 49 was used instead of intermediate 7

NMR (<sup>1</sup>H, CDCl<sub>3</sub>): δ 8.4 (d, 1H), 8.2 (d, 1H), 8.15 (d, 1H), 7.8 (d, 1H), 6.8 (d, 1H), 6.4 (s, 1H), 4.10 (t, 2H), 3.15 (t, 2H), 2.50 (s, 3H), 0.95 (s, 9H), 0.1 (s, 6H).

MS (m/z): 418 [MH]<sup>+</sup>.

Intermediate 51

4-[(4-(1''H-1,3'-Bipyrazol-1'-yl)-3-(2-[(1,1-dimethylethyl)(dimethyl)silyloxy]ethyl)-6-methyl-2-pyridinyl)amino]-3-methylbenzonitrile

As in intermediate 9, except that intermediate 50 was used instead of intermediate 8, and 4-amino-3-methylbenzonitrile was used instead of intermediate 34.

NMR (<sup>1</sup>H, CDCl<sub>3</sub>): δ 8.60 (d, 1H), 8.15 (m, 2H), 7.75 (m, 2H), 7.50 (d, 1H), 7.20 (d, 2H), 6.80 (d, 2H), 6.4 (d, 1H), 4.0 (t, 2H), 3.05 (t, 2H), 2.60 (s, 3H), 2.20 (s, 3H), 0.90 (s, 9H), 0.10 (s, 6H).

MS (m/z): 514 [MH]<sup>+</sup>.

Intermediate 52

4-[(4-(1''H-1,3'-Bipyrazol-1'-yl)-3-(2-[(1,1-dimethylethyl)(dimethyl)silyloxy]ethyl)-6-methyl-2-pyridinyl)amino]-3-(trifluoromethyl)benzonitrile

As in intermediate 9, except that intermediate 50 was used instead of intermediate 8, and 4-amino-3-(trifluoromethyl)benzonitrile was used instead of intermediate 34.

R<sub>f</sub> = 0.25 (7 : 3 CH<sub>2</sub>/EtOAc)

Intermediate 53

4-[(4-(1''H-1,3'-Bipyrazol-1'-yl)-3-(2-[(1,1-dimethylethyl)(dimethyl)silyloxy]ethyl)-6-methyl-2-pyridinyl)amino]-3-chlorobenzonitrile

As in intermediate 9, except that intermediate 50 was used instead of intermediate 8, and 4-amino-3-chlorobenzonitrile was used instead of intermediate 34.

NMR (<sup>1</sup>H, CDCl<sub>3</sub>): δ 8.60 (d, 1H), 8.15 (m, 2H), 7.75-7.65 (m, 2H), 7.50 (d, 1H), 7.20 (d, 2H), 6.80 (d, 2H), 6.4 (d, 1H), 3.90 (t, 2H), 3.15 (t, 2H), 2.20 (s, 3H), 0.90 (s, 9H), 0.10 (s, 6H).

MS (m/z): 535 [MH]<sup>+</sup>.

Intermediate 54

4-[(4-(1''H-1,3'-Bipyrazol-1'-yl)-3-(2-hydroxyethyl)-6-methyl-2-pyridinyl)amino]-3-methylbenzonitrile

As in intermediate 21, except that intermediate 51 was used instead of intermediate 9

NMR (<sup>1</sup>H, CDCl<sub>3</sub>): δ 8.40 (s, 1H), 8.15 (m, 2H), 7.65 (m, 2H), 7.40 (m, 2H), 6.8 (m, 2H), 6.4 (d, 1H), 4.2 (t, 2H), 2.85 (t, 2H), 2.60 (t, 1H), 2.50 (s, 3H), 2.20 (s, 3H).

MS (m/z): 400 [MH]<sup>+</sup>.

Intermediate 55

4-[4-(1'H-1,3'-Bipyrazol-1'-yl)-3-(2-hydroxyethyl)-6-methyl-2-pyridinyl]amino)-3-(trifluoromethyl)benzonitrile

As in intermediate 21, except that intermediate 52 was used instead of intermediate 9

NMR ( $^1\text{H}$ ,  $\text{CDCl}_3$ ):  $\delta$  8.25 (s, 1H), 8.10 (m, 2H), 7.55 (m, 2H), 7.40 (m, 2H), 7.0 (m, 2H), 6.5 (d, 1H), 4.2 (t, 2H), 2.85 (t, 2H), 2.60 (t, 1H), 2.50 (s, 3H).

MS ( $m/z$ ): 454  $[\text{MH}]^+$ .

Intermediate 56

4-[4-(1'H-1,3'-Bipyrazol-1'-yl)-3-(2-hydroxyethyl)-6-methyl-2-pyridinyl]amino)-3-chlorobenzonitrile

As in intermediate 21, except that intermediate 53 was used instead of intermediate 9

NMR ( $^1\text{H}$ ,  $\text{CDCl}_3$ ):  $\delta$  8.35 (s, 1H), 8.15 (m, 2H), 7.65-7.55 (m, 2H), 7.40 (m, 2H), 6.8 (m, 2H), 6.4 (d, 1H), 4.2 (t, 2H), 2.85 (t, 2H), 2.60 (t, 1H), 2.50 (s, 3H).

MS ( $m/z$ ): 420  $[\text{MH}]^+$ .

Intermediate 57

1,2-Dihydro-3H-pyrazol-3-one hydrazone hydrochloride

To a solution of 1H-pyrazol-3-amine (3 g, 1 eq) in 6N HCl (22 mL), at  $-5^\circ\text{C}$ , was added a 1M aqueous solution of  $\text{NaNO}_2$  (36 mL, 1 eq). A solution of  $\text{SnCl}_2$  (13.7 g, 2 eq) in conc. HCl (62 mL) was then added dropwise and the resulting reaction mixture was stirred at r.t. for 2 hr. The solvents were evaporated to give 13 g of a brown solid which was used in the next step without further purification.

Intermediate 58

1'H-1,3'-Bipyrazole

Intermediate 57 was dissolved in a water/ethanol solution (40 mL/28 mL). 1,1,3,3-tetramethoxypropane (8.7 mL, 1.1 eq) was then added and the resulting reaction mixture was stirred for 18 hr at r.t.. The solution was neutralized with  $\text{NaHCO}_3$  and  $\text{K}_2\text{CO}_3$ , extracted with EtOAc (200 mL) and  $\text{CH}_2\text{Cl}_2$  (200 mL). The combined organic extracts were dried over anhyd.  $\text{Na}_2\text{SO}_4$ , the solids were filtered and the solvent evaporated. The crude product was purified by flash chromatography (silica gel,  $\text{CH}_2\text{Cl}_2/\text{EtOAc}$  3:7) to give the title compound as a yellow solid (3.6 g, 74%).

NMR ( $^1\text{H}$ ,  $\text{DMSO}-d_6$ ):  $\delta$  12.45 (s, 1H), 8.25 (bs, 1H), 7.83 (bs, 1H), 7.67 (bs, 1H), 6.45 (bs, 2H).

MS ( $m/z$ ): 134  $[\text{MH}]^+$ .

Intermediate 59

Ethyl 2-chloro-6-methyl-4-[3-(2-pyridinyl)-1H-pyrazol-1-yl]-3-pyridinecarboxylate

As in intermediate 2, except that intermediate 72 (2-(1*H*-pyrazol-3-yl)pyridine) was used instead of 2-(1*H*-pyrazol-3-yl)-1,3-thiazole.

NMR (<sup>1</sup>H, CDCl<sub>3</sub>): δ 8.56 (d, 1H), 7.93 (d, 1H), 7.88 (d, 1H), 7.68 (dt, 1H), 7.17 (m, 2H), 7.11 (d, 1H), 4.30 (q, 2H), 2.54 (s, 3H), 1.18 (t, 3H).

MS (*m/z*): 343 [MH]<sup>+</sup>.

#### Intermediate 60

{2-Chloro-6-methyl-4-[3-(2-pyridinyl)-1*H*-pyrazol-1-yl]-3-pyridinyl}methanol

As in intermediate 3, except that intermediate 59 was used instead of intermediate 2.

NMR (<sup>1</sup>H, CDCl<sub>3</sub>): δ 8.67 (d, 1H), 8.00 (d, 1H), 7.93 (d, 1H), 7.77 (dt, 1H), 7.25-7.19 (m, 3H), 4.77 (d, 2H), 4.44 (t, 1H), 2.59 (s, 3H).

MS (*m/z*): 301 [MH]<sup>+</sup>.

#### Intermediate 61

2-Chloro-6-methyl-4-[3-(2-pyridinyl)-1*H*-pyrazol-1-yl]-3-pyridinecarbaldehyde

As in intermediate 4, except that intermediate 60 was used instead of intermediate 3.

NMR (<sup>1</sup>H, CDCl<sub>3</sub>): δ 10.33 (s, 1H), 8.64 (d, 1H), 8.01 (d, 1H), 7.99 (d, 1H), 7.77 (dt, 1H), 7.34-7.22 (m, 3H), 2.66 (s, 3H).

MS (*m/z*): 299 [MH]<sup>+</sup>.

#### Intermediate 62

2-Chloro-6-methyl-3-[(*E*)-2-(methyloxy)ethenyl]-4-[3-(2-pyridinyl)-1*H*-pyrazol-1-yl]pyridine

As in intermediate 5, except that intermediate 61 was used instead of intermediate 4.

NMR (<sup>1</sup>H, CDCl<sub>3</sub>): δ 8.65 (dd, 1H), 8.02 (dd, 1H), 7.82 (d, 1H), 7.72 (dt, 1H), 7.36 (s, 1H), 7.26 (m, 1H), 7.09 (d, 1H), 6.48 (d, 1H), 5.61 (d, 1H), 3.60 (s, 3H), 2.54 (s, 3H).

MS (*m/z*): 327 [MH]<sup>+</sup>.

#### Intermediate 63

{2-Chloro-6-methyl-4-[3-(2-pyridinyl)-1*H*-pyrazol-1-yl]-3-pyridinyl}acetaldehyde

As in intermediate 6, except that intermediate 62 was used instead of intermediate 5.

NMR (<sup>1</sup>H, CDCl<sub>3</sub>): δ 9.82 (s, 1H), 8.62 (d, 1H), 7.92 (d, 1H), 7.81 (d, 1H), 7.74 (dt, 1H), 7.27-7.16 (m, 3H), 4.08 (s, 2H), 2.60 (s, 3H).

MS (*m/z*): 313 [MH]<sup>+</sup>.

#### Intermediate 64

2-{2-Chloro-6-methyl-4-[3-(2-pyridinyl)-1*H*-pyrazol-1-yl]-3-pyridinyl}ethanol

As in intermediate 7, except that intermediate 63 was used instead of intermediate 6.

NMR (<sup>1</sup>H, CDCl<sub>3</sub>): δ 8.65 (d, 1H), 7.95 (dd, 1H), 7.82 (d, 1H), 7.76 (dt, 1H), 7.27 (m, 1H), 7.19 (d, 1H), 7.09 (s, 1H), 5.07 (bs, 1H), 4.11 (t, 2H), 3.09 (t, 2H), 2.57 (s, 3H).

MS (*m/z*): 315 [MH]<sup>+</sup>.

Intermediate 65

2-chloro-3-(2-((1,1-dimethylethyl)(dimethyl)silyloxy)ethyl)-6-methyl-4-[3-(2-pyridinyl)-1H-pyrazol-1-yl]pyridine

As in intermediate 8, except that intermediate 64 was used instead of intermediate 7.

NMR (<sup>1</sup>H, CDCl<sub>3</sub>): δ 8.64 (dd, 1H), 8.17 (dd, 1H), 8.04 (d, 1H), 7.72 (dt, 1H), 7.24 (m, 2H), 7.12 (d, 1H), 3.99 (t, 2H), 3.10 (t, 2H), 2.55 (s, 3H), 0.79 (s, 9H), 0.05 (s, 6H).

MS (m/z): 429 [MH]<sup>+</sup>.

Intermediate 66

4-((3-(2-((1,1-Dimethylethyl)(dimethyl)silyloxy)ethyl)-6-methyl-4-[3-(2-pyridinyl)-1H-pyrazol-1-yl]-2-pyridinyl)amino)-3-methylbenzonitrile

As in intermediate 9, except that intermediate 65 was used instead of intermediate 8, and 4-amino-3-methylbenzonitrile was used instead of intermediate 34.

NMR (<sup>1</sup>H, CDCl<sub>3</sub>): δ 8.64-6.59 (m, 7H), 4.23 (t, 2H), 4.07 (bs, 1H), 2.83 (t, 2H), 2.53 (s, 3H), 2.32 (s, 3H), 0.81 (s, 9H), 0.03 (s, 6H).

MS (m/z): 525 [MH]<sup>+</sup>.

Intermediate 67

3-Chloro-4-((3-(2-((1,1-dimethylethyl)(dimethyl)silyloxy)ethyl)-6-methyl-4-[3-(2-pyridinyl)-1H-pyrazol-1-yl]-2-pyridinyl)amino)benzonitrile

As in intermediate 9, except that intermediate 65 was used instead of intermediate 8, and 4-amino-3-chlorobenzonitrile was used instead of intermediate 34.

NMR (<sup>1</sup>H, CDCl<sub>3</sub>): δ 8.64 (d, 1H), 8.59 (d, 1H), 8.17 (s, 1H), 8.05 (d, 1H), 7.75 (d, 1H), 7.62 (s, 1H), 7.62 (s, 1H), 7.49 (d, 1H), 7.22 (s, 1H), 7.12 (s, 1H), 6.88 (s, 1H), 3.99 (t, 2H), 3.10 (t, 2H), 2.55 (s, 3H), 0.79 (s, 9H), 0.05 (s, 6H).

MS (m/z): 545 [MH]<sup>+</sup>.

Intermediate 68

4-((3-(2-((1,1-Dimethylethyl)(dimethyl)silyloxy)ethyl)-6-methyl-4-[3-(2-pyridinyl)-1H-pyrazol-1-yl]-2-pyridinyl)amino)-3-(trifluoromethyl)benzonitrile

As in intermediate 9, except that intermediate 65 was used instead of intermediate 8, and 4-amino-3-(trifluoromethyl)benzonitrile was used instead of intermediate 34.

NMR (<sup>1</sup>H, CDCl<sub>3</sub>): δ 8.64-6.93 (m, 7H), 4.7 (bs, 1H), 4.10 (t, 2H), 2.88 (t, 2H), 2.55 (s, 3H), 0.77 (s, 9H), 0.08 (s, 6H).

MS (m/z): 579 [MH]<sup>+</sup>.

Intermediate 69

4-((3-(2-Hydroxyethyl)-6-methyl-4-[3-(2-pyridinyl)-1H-pyrazol-1-yl]-2-pyridinyl)amino)-3-methylbenzonitrile

As in intermediate 21, except that intermediate 66 was used instead of intermediate 9.

NMR ( $^1\text{H}$ ,  $\text{CDCl}_3$ ):  $\delta$  8.65 (dd, 1H), 8.15 (m, 2H), 7.95 (d, 1H), 7.75 (m, 2H), 7.42 (m, 2H), 7.26 (m, 1H), 7.15 (d, 1H), 6.80 (s, 1H), 4.27 (t, 2H), 2.84 (t, 2H), 2.51 (s, 3H), 2.30 (s, 3H).  
MS ( $m/z$ ): 411  $[\text{MH}]^+$ .

#### Intermediate 70

##### 3-Chloro-4-((3-(2-hydroxyethyl)-6-methyl-4-[3-(2-pyridinyl)-1H-pyrazol-1-yl]-2-pyridinyl)amino)benzonitrile

As in intermediate 21, except that intermediate 67 was used instead of intermediate 9.

NMR ( $^1\text{H}$ ,  $\text{CDCl}_3$ ):  $\delta$  8.65 (d, 1H), 8.55 (d, 1H), 8.46 (s, 1H), 7.96 (d, 1H), 7.77 (m, 2H), 7.64 (s, 1H), 7.50 (d, 1H), 7.26 (d, 1H), 7.17 (d, 1H), 6.85 (s, 1H), 4.25 (t, 2H), 3.72 (bs, 1H), 2.92 (t, 2H), 2.54 (s, 3H).

MS ( $m/z$ ): 431  $[\text{MH}]^+$ .

#### Intermediate 71

##### 4-((3-(2-Hydroxyethyl)-6-methyl-4-[3-(2-pyridinyl)-1H-pyrazol-1-yl]-2-pyridinyl)amino)-3-(trifluoromethyl)benzonitrile

As in intermediate 21, except that intermediate 68 was used instead of intermediate 9.

NMR ( $^1\text{H}$ ,  $\text{CDCl}_3$ ):  $\delta$  8.65 (dd, 1H), 8.53 (bs, 1H), 8.31 (d, 1H), 7.95 (d, 1H), 7.84 (dd, 1H), 7.77-7.68 (m, 3H), 7.25 (m, 1H), 7.17 (d, 1H), 6.80 (s, 1H), 4.18 (m, 2H), 3.41 (bs, 1H), 2.87 (t, 2H), 2.52 (s, 3H).

MS ( $m/z$ ): 465  $[\text{MH}]^+$ .

#### Intermediate 72

##### 2-(1H-Pyrazol-3-yl)pyridine

The title compound was prepared according to an already published procedure: Pleier, Anna-Katharina; Glas, Holger; Grosche, Manja; Sirsch, Peter; Thiel, Werner R.; *Synthesis*, 2001, 1, p.55-62

#### Intermediate 73

##### 2-Methyl-6-(methoxy)pyridine

To a solution of 6-methyl-pyridin-2-ol (2.5 g, 0.023 mol) in  $\text{CH}_2\text{Cl}_2$  (10 mL) at r.t., under  $\text{N}_2$ , were added  $\text{Ag}_2\text{CO}_3$  (6 g, 1.5 eq.) and MeI (5.6 mL, 4 eq). The solution was stirred at r.t. for 4 days, then  $\text{Ag}_2\text{CO}_3$  was filtered and washed with  $\text{CH}_2\text{Cl}_2$ , and the organic layer was evaporated to dryness. The crude product was purified by flash chromatography (silica gel, EtOAc/hex 2:8) to give the title compound (1.9 mg, 67%) as a white solid.

NMR ( $^1\text{H}$ ,  $\text{CDCl}_3$ ):  $\delta$  7.4 (t, 1H), 6.6 (d, 1H), 6.5 (d, 1H), 3.8 (s, 3H), 2.4 (s, 3H).

#### Intermediate 74

##### 2-Methyl-6-(methoxy)-3-nitropyridine

To intermediate 73 (1.95 g, 0.015 mol) at r.t., was added  $\text{HNO}_3$  60% (7ml). The mixture became reddish brown with generation of heat and  $\text{NO}_2$ . Conc.  $\text{H}_2\text{SO}_4$  (7ml) was then added.

The solution was stirred at 80°C overnight. Into the cooled reaction mixture was poured ice water (70 mL) and the solution was neutralized with CaCO<sub>3</sub>. The aqueous solution was extracted with EtOAc (3x50 mL) and the combined organic extracts were dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>. The solids were filtered and the solvent was evaporated to dryness to give the title compound (2.25 g, 86%) as a yellow solid.

NMR (<sup>1</sup>H, DMSO): δ 8.3 (d, 1H); 6.8 (d, 1H); 4 (s, 3H), 2.7 (s, 3H).

#### Intermediate 75

##### 2-Methyl-6-(methoxy)-3-pyridinamine

To a solution of Intermediate 74 (2 g, 0.012 mol) in H<sub>2</sub>O/MeOH 1:1 (40 mL), were added NH<sub>4</sub>Cl (2.2 g, 3.5 eq.) and Fe powder (2.3 g, 3.5 eq.). The reaction mixture was stirred at 80°C for 16 hr. Fe was then filtered and washed with MeOH. The MeOH was evaporated, H<sub>2</sub>O was added and the aqueous solution was extracted with EtOAc (3x50 mL). The combined organic extracts were dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, the solids were filtered and the solvent evaporated to dryness to give the title compound (1.35 g, 81%) as a brown oil.

NMR (<sup>1</sup>H, CDCl<sub>3</sub>): δ 6.9 (d, 1H), 6.4 (d, 1H), 3.8 (s, 3H), 2.33 (s, 3H).

#### Intermediate 76

##### {2-Chloro-6-methyl-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-3-pyridinyl}methyl methanesulfonate

To a solution of Intermediate 3 (1 g, 3.26 mmols) in anhydrous CH<sub>2</sub>Cl<sub>2</sub>/DMF (15/15 mL), at -15°C, under N<sub>2</sub>, were slowly added Et<sub>3</sub>N (0.92 mL, 2 eq) and MsCl (0.39 mL, 1.5 eq). The reaction mixture was stirred at -15°C for 2.5 hr. It was then quenched with water (10 mL) and extracted with CH<sub>2</sub>Cl<sub>2</sub> (3x20 mL). The combined organic extracts were washed with sat. aq. NaCl (10 mL) and dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>. The solids were filtered and the solvent evaporated. The crude product was purified by flash chromatography (silica gel, cHex/EtOAc 6:4 → 4:6) to give the title compound as a white solid (1 g, 85%).

NMR (<sup>1</sup>H, CDCl<sub>3</sub>): δ 7.90 (d, 1H), 7.87 (d, 1H), 7.39 (d, 1H), 7.34 (s, 1H), 7.14 (d, 1H), 5.5 (s, 2H), 3.00 (s, 3H), 2.78 (s, 3H).

MS (m/z): 385 [MH]<sup>+</sup>.

#### Intermediate 77

##### {2-Chloro-6-methyl-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-3-pyridinyl}acetonitrile

To a solution of Intermediate 76 (8 g, 20.79 mmols) in anhydrous DMF (200 mL), at 0°C, under N<sub>2</sub>, was added KCN (1.35 g, 1 eq). The reaction mixture was stirred at r.t. for 6 hr. It was then diluted with Et<sub>2</sub>O (500 mL), washed with 1M NaOH (2x200 mL), with sat. aq. NaCl (2x200 mL) and dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>. The solids were filtered and the solvent evaporated. The title compound was obtained as a pale brown solid (6.5 g, 96%) and was used in the next step without further purification.

NMR (<sup>1</sup>H, CDCl<sub>3</sub>): δ 7.92 (d, 1H), 7.91 (d, 1H), 7.41 (d, 1H), 7.31 (s, 1H), 7.18 (d, 1H), 3.99 (s, 2H), 2.78 (s, 3H).

MS (*m/z*): 316 [MH]<sup>+</sup>.

Intermediate 78

(2-{2-Chloro-6-methyl-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-3-pyridinyl}ethyl)amine hydrochloride

To a solution of Intermediate 77 (6 g, 19.0 mmols) in anh. THF (6 mL), at r.t., under N<sub>2</sub>, was added BH<sub>3</sub>·THF 1M/THF (3 eq., 57 mL). The reaction mixture was stirred at reflux for 2 hr. It was then cooled to r.t. and HCl 1M/Et<sub>2</sub>O (6 eq with respect to BH<sub>3</sub>·THF, exothermic reaction) and MeOH (10 mL) were slowly added. The resulting solution was heated at reflux for 30 min and then concentrated *in vacuo*. The crude product was dissolved in absolute EtOH and cooled to 0°C. Et<sub>2</sub>O was added (60 mL) and a precipitate formed. The solid was filtered and dried to give the title compound as a pale yellow solid (6.7 g, 98%).

NMR (<sup>1</sup>H, CDCl<sub>3</sub>): δ 8.42 (d, 1H), 7.94 (d, 1H), 7.79 (d, 1H), 7.48 (s, 1H), 7.07 (d, 1H), 2.81 (m, 4H), 2.51 (s, 3H), 2.0 (bs, 2H).

MS (*m/z*): 320.3 [MH]<sup>+</sup>.

Intermediate 79

6-Methyl-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine

To a solution of Intermediate 78 (1 g, 2.81 mmols) in anh. NMP (15 mL), at r.t., under N<sub>2</sub>, were added CuI (107 mg, 0.2 eq), K<sub>2</sub>CO<sub>3</sub> (1.17 g, 3 eq) and (1R,2R)-diaminomethylcyclohexane (240 mg, 0.6 eq). The reaction mixture was stirred at 150°C for 4 hr. It was then diluted with water and extracted with EtOAc (2x20 mL). The combined organic extracts were washed with sat. aq. NaCl and dried over anh. Na<sub>2</sub>SO<sub>4</sub>. The solids were filtered and the solvent evaporated *in vacuo*. The crude product was purified by flash chromatography (silica gel CH<sub>2</sub>Cl<sub>2</sub>/MeOH 98:2). The title compound was obtained as a white solid (400 mg, 50%).

NMR (<sup>1</sup>H, CDCl<sub>3</sub>): δ 7.98 (d, 1H), 7.89 (d, 1H), 7.35 (d, 1H), 7.06 (d, 1H), 4.65 (bs, 1H), 3.72 (t, 2H), 3.48 (t, 2H), 2.42 (s, 3H)

MS (*m/z*): 284.3 [MH]<sup>+</sup>

Intermediate 80

1-[2-Methyl-4-(methyloxy)phenyl]-2-pyrrolidinone

To a solution of Et<sub>3</sub>N (156 mL, 1 eq) and 2-methyl-4-(methyloxy)aniline (150 g, 1.09 mole) in anh. THF (2.4 L), in a 10 L reaction vessel, at 0°C, under N<sub>2</sub>, was added dropwise a solution of 4-chlorobutyl chloride (126 mL, 1 eq) in anh. THF (480 mL). The internal temperature was maintained at *circa* 10°C and the reaction mixture was stirred for 1.5 hr. It was cooled down to 0°C and KOt-Bu 1M/THF (2.64 L, 2.4 eq) was added dropwise over a period of 1.5 hr, keeping the internal temperature <10°C. The reaction mixture was stirred at that temperature for 30 min. Water (1.5 L) was then added slowly (20 min) and the phases were separated. The organic layer was treated with conc. HCl (250 mL) and water (1.26 L) and the phases were separated. The combined aqueous layers were extracted with EtOAc (2.6 L)

and the combined organic layers were washed with sat.aq. NaCl (2 L). The solvent was evaporated and the residue purified by flash chromatography (Biotage 150, EtOAc/cHex 8:2) to give the title compound as a pale brown solid (206 g, 92%).

NMR ( $^1\text{H}$ ,  $\text{CDCl}_3$ ):  $\delta$  7.05 (d, 1H), 6.79-6.72 (m, 2H), 3.75 (s, 3H), 3.64 (t, 2H), 2.18 (s, 6H).

MS (m/z): 206 [MH] $^+$ .

#### Intermediate 81

##### Ethyl 3-((1-[2-methyl-4-(methyloxy)phenyl]-2-pyrrolidinylidene)amino)-2-butenate

To a solution of intermediate 80 (8.3 g, 40.49 mmols) in anh. 1,2-dichloroethane (100 mL), at r.t., under  $\text{N}_2$ , was added  $\text{POCl}_3$  (7.5 mL, 2 eq) dropwise followed by ethyl 3-aminocrotonate (5.17 mL, 1 eq). The reaction mixture was heated at  $60^\circ\text{C}$  for 3.5 hr. It was then cooled down to r.t. and neutralized to pH 7 by the careful addition of sat.aq.  $\text{NaHCO}_3$ . The neutralized solution was extracted with  $\text{CH}_2\text{Cl}_2$  (3x50 mL). The combined organic extracts were washed with sat.aq. NaCl and dried over anh.  $\text{Na}_2\text{SO}_4$ . The solids were filtered and the solvent evaporated. The crude product was used as such in the next step (17.8 g).

MS (m/z): 317 [MH] $^+$ .

#### Intermediate 82

##### 6-Methyl-1-[2-methyl-4-(methyloxy)phenyl]-1,2,3,7-tetrahydro-4H-pyrrolo[2,3-b]pyridin-4-one

A solution of intermediate 81 (17.8 g, 55 mmols) in anh. DMF (50 mL) was added dropwise to a suspension of NaH 60%/oil (4.5 g, 2 eq) in anh. DMF. The reaction mixture was heated at  $100^\circ\text{C}$  for 8 hr. More NaH 60%/oil (2.25 g, 1 eq) was added and the reaction mixture was heated for an additional 4 hr. It was cooled down to r.t. and carefully poured in sat.aq.  $\text{NH}_4\text{Cl}$ . The aqueous solution was extracted with  $\text{CH}_2\text{Cl}_2$  (5x50 mL) and the combined organic extracts were dried over anh.  $\text{Na}_2\text{SO}_4$ . The solids were filtered and the solvent was evaporated. The crude compound was purified by flash chromatography (Biotage 75,  $\text{CH}_2\text{Cl}_2/\text{MeOH}$  95:5  $\rightarrow$  80:20). The title compound was obtained as a brown oil (952 mg, 9%, two steps)

NMR ( $^1\text{H}$ ,  $\text{CDCl}_3$ ):  $\delta$  7.08 (d, 1H), 6.72-6.68 (m, 2H), 5.87 (s, 1H), 3.73 (s, 3H), 3.73 (t, 2H), 2.99 (t, 2H), 2.21 (s, 3H), 2.13 (s, 3H).

MS (m/z): 271 [MH] $^+$ .

#### Intermediate 83

##### 6-Methyl-1-[2-methyl-4-(methyloxy)phenyl]-2,3-dihydro-1H-pyrrolo[2,3-b]pyridin-4-yl trifluoromethanesulfonate

To a solution of intermediate 82 (950 mg, 3.52 mmols) in anh.  $\text{CH}_2\text{Cl}_2$  (10 mL), at  $-20^\circ\text{C}$ , under  $\text{N}_2$ , were added pyridine (626  $\mu\text{L}$ , 2.2 eq) and triflic anhydride (651  $\mu\text{L}$ , 1.1 eq). The reaction mixture was stirred at r.t. for 2 hr. It was then poured in sat.aq.  $\text{NH}_4\text{Cl}$  (20 mL) and the phases were separated. The aqueous layer was extracted with  $\text{CH}_2\text{Cl}_2$  (2x10 mL) and the combined organic extracts were dried over anh.  $\text{Na}_2\text{SO}_4$ . The solids were filtered and the



solvent evaporated. The crude product was purified by flash chromatography (silica gel, cHex/EtOAc 9:1) to give the title compound as a white solid (913 mg, 64%).

NMR ( $^1\text{H}$ ,  $\text{CDCl}_3$ ):  $\delta$  7.12 (d, 1H), 6.81-6.75 (m, 2H), 6.24 (s, 1H), 3.89 (t, 2H), 3.80 (s, 3H), 3.21 (t, 2H), 2.29 (s, 3H), 2.21 (s, 3H).

MS ( $m/z$ ): 403  $[\text{MH}]^+$ .

#### Intermediate 84

##### 4-Iodo-6-methyl-1-[2-methyl-4-(methoxy)phenyl]-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine

To a solution of intermediate 83 (913 mg, 2.27 mmols) in anhyd. NMP (7 mL), at r.t., under  $\text{N}_2$ , was added KI (1.13 g, 3 eq) and the reaction mixture was stirred at  $150^\circ\text{C}$  for 18 hr. It was then cooled down to r.t. and diluted in water/sat. aq. NaCl. The aqueous phase was extracted with EtOAc (3x30 mL) and the combined organic extracts were dried over anhyd.  $\text{Na}_2\text{SO}_4$ . The solids were filtered and the solvent evaporated. The crude product was purified by flash chromatography (silica gel, cHex/EtOAc 9:1) to give the title compound as a clear oil, which solidified upon standing (681 mg, 79%).

NMR ( $^1\text{H}$ ,  $\text{CDCl}_3$ ):  $\delta$  7.14 (d, 1H), 6.81-6.74 (m, 2H), 6.70 (s, 1H), 3.84 (t, 2H), 3.81 (s, 3H), 3.03 (t, 2H), 2.22 (s, 6H).

MS ( $m/z$ ): 381  $[\text{MH}]^+$ .

#### EXAMPLE 1

##### Synthesis of representative compounds of structure (IVa)

#### Example 1-1

##### 6-Methyl-1-[6-(methoxy)-2-(trifluoromethyl)-3-pyridinyl]-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine

To a solution of intermediate 21 (10 mg, 0.021 mmol) in anhyd. DMF (1.5 mL) at r.t., were added  $\text{Et}_3\text{N}$  (12  $\mu\text{L}$ , 4 eq) and  $\text{MsCl}$  (4  $\mu\text{L}$ , 2 eq). The solution was stirred at  $56^\circ\text{C}$  for 4 hr.  $\text{CH}_2\text{Cl}_2$  (5 mL) was added and the reaction quenched with  $\text{H}_2\text{O}$  (3 mL). The aqueous phase was extracted with  $\text{CH}_2\text{Cl}_2$  (3x2 mL) and the combined organic extracts dried over anhyd.  $\text{Na}_2\text{SO}_4$ . The solids were filtered and the solvent evaporated to dryness. The crude product was purified by flash chromatography (silica gel,  $\text{CH}_2\text{Cl}_2/\text{MeOH}$  99:1) to give the title compound (2 mg, 21 %) as a white foam.

#### Example 1-2

##### 6-Methyl-1-[2-methyl-6-(methoxy)-3-pyridinyl]-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine

To a solution of intermediate 22 (126 mg, 0.030 mmol) in anhyd.  $\text{CH}_2\text{Cl}_2$  (10 mL) at r.t., under  $\text{N}_2$ , were added  $\text{Et}_3\text{N}$  (0.0167 mL, 4eq),  $\text{PPh}_3$  (310 mg, 4eq) and  $\text{I}_2$  (305 mg, 4eq). The reaction mixture was stirred at r.t. for 2 hr. Water (10 mL) was then added and the solution extracted with EtOAc (15 mL). The organic layer was dried over anhyd.  $\text{Na}_2\text{SO}_4$ , the solids were filtered and the solvent evaporated *in vacuo*. The crude compound thus obtained was

purified by flash chromatography (silica gel, cHex/EtOAc 6:4) to give 5.9 mg of the title compound as a white solid.

#### Example 1-3

##### 1-[2,6-Bis(methoxy)-3-pyridinyl]-6-methyl-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine


To a solution of intermediate 23 (46mg, 0.105 mmol) in anh.  $\text{CH}_2\text{Cl}_2$  (2 mL) at r.t., under  $\text{N}_2$ , were added  $\text{Et}_3\text{N}$  (73  $\mu\text{L}$ , 5eq), polymer bound  $\text{PPh}_3$  (174mg, 5eq) and  $\text{CBr}_4$  (174mg, 5eq). The reaction mixture was stirred at r.t. for 2 hr. Water (10 mL) was then added and the solution extracted with EtOAc (15 mL). The organic layer was dried over anh.  $\text{Na}_2\text{SO}_4$ , the solids were filtered and the solvent evaporated *in vacuo*. The crude compound thus obtained was purified by flash chromatography (silica gel, cHex/EtOAc 7:3) to give 14 mg of the title compound as a white solid.

#### Example 1-4

##### N,N,4-Trimethyl-5-[6-methyl-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-2,3-dihydro-1H-pyrrolo[2,3-b]pyridin-1-yl]-2-pyridinamine

To a solution of intermediate 24 (45 mg, 0.1 mmol) in anh.  $\text{CH}_2\text{Cl}_2$  (2 mL), at r.t., under  $\text{N}_2$ , were added  $\text{I}_2$  (50.8 mg, 2 eq),  $\text{PPh}_3$  (52.4 mg, 2 eq) and  $\text{Et}_3\text{N}$  (27  $\mu\text{L}$ , 2 eq). The reaction mixture was stirred at r.t. for 2.5 hr and evaporated to dryness. The crude product was purified by flash chromatography (silica gel, EtOAc/MeOH 95:5  $\rightarrow$  7:3 EtOAc/ $\text{NH}_3$  (0.5 M in MeOH)) to give the title compound as a yellow solid (3 mg, 10 %).

All the analytical data are set forth in the following Table 1-1.



Cpd. No.	W	R <sub>1</sub>	Analytical Data
1-1	2-trifluoromethyl-4-methoxypyridin-3-yl	CH <sub>3</sub>	NMR ( <sup>1</sup> H, CDCl <sub>3</sub> ): δ 7.9 (d, 1H); 7.8 (d, 1H); 7.6 (d, 1H); 7.3 (d, 1H); 7.0 (d, 1H); 6.9 (d, 1H); 6.7 (s, 1H); 3.9 (s, 3H); 3.8 (t, 2H); 3.5 (t, 2H); 2.33 (s, 3H). MS (m/z): 459 [MH] <sup>+</sup> .
1-2	2-methyl-4-methoxy-pyridin-3-yl	CH <sub>3</sub>	NMR ( <sup>1</sup> H, CDCl <sub>3</sub> ): δ 7.98 (d, 1H), 7.88 (d, 1H), 6.98 (d, 1H), 6.83 (d, 1H), 7.03 (d, 1H), 6.62 (d, 1H), 6.60 (s, 1H), 3.92 (s, 3H), 3.90 (d, 2H), 3.56 (d, 2H), 2.39 (s, 3H), 2.35 (s, 3H). MS (m/z): 405 [MH] <sup>+</sup> .
1-3	2,4-dimethoxypyridin-3-yl	CH <sub>3</sub>	NMR ( <sup>1</sup> H, CDCl <sub>3</sub> ): δ 7.98 (d, 1H); 7.87 (d, 1H); 7.72 (d, 1H); 7.34 (d, 1H); 7.06 (d, 1H); 6.67 (s, 1H); 6.36 (d, 1H); 3.96 (t, 2H); 3.96 (s, 3H); 3.94 (s, 3H); 3.51 (t, 2H); 2.37 (s, 3H)δ. MS (m/z): 421 [MH] <sup>+</sup> .
1-4	6-methyl-4-dimethyl-amino-pyridin-3-yl	CH <sub>3</sub>	NMR ( <sup>1</sup> H, DMSO-d <sub>6</sub> ): δ 8.57 (d, 1H), 7.95 (s, 1H), 7.92 (d, 1H), 7.78 (d, 1H), 7.04 (d, 1H), 6.86 (s, 1H), 6.55 (s, 1H), 3.88 (t, 2H), 3.49 (t, 2H), 3.16 (s, 3H), 3.14 (s, 3H), 3.01 (s, 6H). IR (, cm <sup>-1</sup> ): MS (m/z): 418 [MH] <sup>+</sup> .

## EXAMPLE 2

## Synthesis of representative compounds of structure (Va)

Example 2-11-(2-Difluoromethyl-4-methoxy-phenyl)-6-methyl-4-(3-thiazol-2-yl-pyrazol-1-yl)-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine

To a solution of Intermediate 25 (42 mg, 0.092 mmol) in anh. CH<sub>2</sub>Cl<sub>2</sub> (8 mL) at r.t., under N<sub>2</sub>, were added PPh<sub>3</sub> (48 mg, 2 eq) and CBr<sub>4</sub> (61 mg, 2 eq). The reaction mixture was stirred at r.t. for 6 hr. Sat.aq. NaHCO<sub>3</sub> (10 mL) was added. The phases were separated and the aqueous layer was extracted with EtOAc (2x20mL) and the combined organic extracts were dried over anh. Na<sub>2</sub>SO<sub>4</sub>. The solids were filtered and the solvent was evaporated to dryness.

The crude product was purified by flash chromatography (silica gel, cHex/EtOAc 8:2) to give the title compound (19 mg, 50 %) as a white solid.

#### Example 2-2

##### 1-(2-Chloro-4-methoxy-phenyl)-6-methyl-4-[3-(1,3-thiazol-2-yl)-pyrazol-1-yl]-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine

To a solution of intermediate 26 (25 mg, 0.056 mmol) in anh. DMF (2 mL) at r.t., under N<sub>2</sub>, were added Et<sub>3</sub>N (30 µL, 4 eq) and MsCl (10 µL, 2 eq). The solution was stirred at 50°C for 5 hr then CH<sub>2</sub>Cl<sub>2</sub> (5 mL) was added and the reaction quenched with H<sub>2</sub>O (2 mL). The phases were separated and the aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3x3 mL). The combined organic extracts were dried over anh. Na<sub>2</sub>SO<sub>4</sub>, the solids were filtered and the solvent evaporated under reduced pressure. The crude product was purified by flash chromatography (silica gel, CH<sub>2</sub>Cl<sub>2</sub>/MeOH 99:1) to give the title compound (8 mg, 33 %) as a white foam.

#### Example 2-3

##### 1-[2,4-Bis(methoxy)phenyl]-6-methyl-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine

To a suspension of intermediate 27 (26 mg, 0.06 mmol) in anh. CH<sub>2</sub>Cl<sub>2</sub> (1 mL), at r.t., under N<sub>2</sub>, was added Et<sub>3</sub>N (17 µL, 3 eq), PPh<sub>3</sub> (31 mg, 2 eq) and I<sub>2</sub> (3 mg, 2 eq). The reaction mixture was stirred at r.t. for 24 hr and then partitioned between CH<sub>2</sub>Cl<sub>2</sub>/H<sub>2</sub>O. The phases were separated and the organic layer was dried over anh. Na<sub>2</sub>SO<sub>4</sub>. The solids were filtered and the solvent evaporated. The crude product was purified by flash chromatography (silica gel, 100% CH<sub>2</sub>Cl<sub>2</sub> → CH<sub>2</sub>Cl<sub>2</sub>/MeOH 97:3) to give the title compound as a white solid (5 mg, 20%).

#### Example 2-4

##### 3-Chloro-4-[6-methyl-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-2,3-dihydro-1H-pyrrolo[2,3-b]pyridin-1-yl]benzonitrile

To a solution of intermediate 79 (50 mg, 0.176 mmol) in anh. DME (1 mL) were added Pd<sub>2</sub>(dba)<sub>3</sub> (16 mg, 0.1 eq), 2-(dicyclohexylphosphino)-2'-methylbiphenyl (19 mg, 0.3 eq), K<sub>3</sub>PO<sub>4</sub> (101 mg, 2.7 eq) and 4-amino-3-chlorobenzonitrile (47 mg, 1 eq). The reaction mixture was subjected to microwave irradiation (150W, 100°C, 60 psi) for 20 min. It was then quenched with sat.aq. NH<sub>4</sub>Cl and extracted with EtOAc (2x10 mL). The combined organic extracts were washed with sat.aq. NaCl and dried over anh. Na<sub>2</sub>SO<sub>4</sub>. The solids were filtered and the solvent evaporated. The crude product was purified by flash chromatography (silica gel, cHex/EtOAc 8:2 → 7:3) to give the title compound as a white solid (16 mg, 22%).

#### Example 2-5

##### 4-[6-Methyl-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-2,3-dihydro-1H-pyrrolo[2,3-b]pyridin-1-yl]-3-[(trifluoromethyl)oxy]benzonitrile

To a solution of intermediate 28 (93 mg, 0.191 mmol) in anh. CH<sub>2</sub>Cl<sub>2</sub> (3 mL), at r.t., under N<sub>2</sub>, were added CBr<sub>4</sub> (317 mg, 5eq) and PPh<sub>3</sub> (317 mg, 5eq). The reaction mixture was stirred at r.t. for 2 hr. Water (5 mL) was then added and the solution extracted with CH<sub>2</sub>Cl<sub>2</sub> (2x10 mL). The combined organic extracts were dried over anh. Na<sub>2</sub>SO<sub>4</sub>, the solids were filtered and the solvents evaporated *in vacuo*. The crude compound thus obtained was purified by flash chromatography (silica gel, cHex/EtOAc 1:1) to give 37 mg of the title compound as a white solid.

#### Example 2-6

##### 3-Ethyl-4-[6-methyl-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-2,3-dihydro-1H-pyrrolo[2,3-b]pyridin-1-yl]benzonitrile

To a suspension of intermediate 29 (45 mg, 0.105 mmol) in anh. CH<sub>2</sub>Cl<sub>2</sub> (2 mL), at r.t., under N<sub>2</sub>, was added Et<sub>3</sub>N (58 µL, 4 eq) and MsCl (16 µL, 2 eq). The reaction mixture was stirred at r.t. for 24 hr and then partitioned between CH<sub>2</sub>Cl<sub>2</sub>/H<sub>2</sub>O. The phases were separated and the organic layer was dried over anh. Na<sub>2</sub>SO<sub>4</sub>. The solids were filtered and the solvent evaporated. The crude product was purified by flash chromatography (silica gel, 100% CH<sub>2</sub>Cl<sub>2</sub> → CH<sub>2</sub>Cl<sub>2</sub>/MeOH 98:2) to give the title compound as a white solid (2 mg, 5%).

#### Example 2-7

##### 1-(4-Fluoro-2-methylphenyl)-6-methyl-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine

To a solution of intermediate 30 (7.6 mg, 1 eq) in anh. CH<sub>2</sub>Cl<sub>2</sub> (5 mL), at r.t., under N<sub>2</sub>, were added Et<sub>3</sub>N (9 µL, 3.5 eq) and methanesulfonyl chloride (4.3 µL, 3 eq) and the reaction mixture was stirred at r.t. for 2 hr. It was then partitioned between EtOAc/sat.aq. NaCl. The phases were separated and the organic layer was washed with sat.aq. NaCl (2x10 mL). It was dried over anh. Na<sub>2</sub>SO<sub>4</sub>, the solids were filtered and the solvent evaporated. The crude product was purified by flash chromatography (silica gel, cHex/EtOAc 7:3) to give the title compound as a yellow solid (0.9 mg, 13%).


#### Example 2-8

##### 6-Methyl-1-[2-methyl-4-(1H-pyrazol-1-yl)phenyl]-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine

To a solution of intermediate 31 (30 mg, 0.065 mmol) in anh. CH<sub>2</sub>Cl<sub>2</sub> (2 mL), at r.t., under N<sub>2</sub>, were added I<sub>2</sub> (66 mg, 4 eq), PPh<sub>3</sub> (68 mg, 4 eq) and Et<sub>3</sub>N (36 µL, 4 eq). The reaction mixture was stirred at r.t. for 3.5 hr. Sat.aq. NaCl was added and the phases were separated. The aqueous layer was further extracted with CH<sub>2</sub>Cl<sub>2</sub> (2x10 mL) and EtOAc (2x10 mL). The combined organic extracts were dried over anh. Na<sub>2</sub>SO<sub>4</sub>, the solids were filtered and the solvent evaporated. The residue was purified by SCX Column (Eluents: CH<sub>2</sub>Cl<sub>2</sub>, MeOH and NH<sub>3</sub>(0.5 M in MeOH)), then by preparative HPLC (Column: X Terra MS C18 5 µm, 100 x 19 mm / Mobile phase: A: H<sub>2</sub>O + 0.1% TFA; B: CH<sub>3</sub>CN + 0.1% TFA / Gradient: 25% (B) for 6 min, from 25% (B) to 50% (B) in 10 min / Flow rate: 17 mL/min / UV wavelength range: 200-

400 nm / Mass range: 100-900 amu / Ionization: ES+) to give the title compound as a white solid (5 mg, 17%).

All the analytical data are set forth in the following Table 2-1.



Cpd. No.	W1	R <sub>1</sub>	Analytical Data
2-1	2-difluoromethyl-4-methoxyphenyl	CH <sub>3</sub>	NMR ( <sup>1</sup> H, CDCl <sub>3</sub> ): δ 8.00 (d, 1H), 7.89 (d, 1H), 7.36 (d, 1H), 7.24 (m, 2H), 7.08 (m, 2H), 6.88 (t, 1H; J <sub>0+H</sub> = 55.8 Hz), 6.71 (s, 1H), 3.95 (t, 2H), 3.87 (s, 3H), 3.56 (t, 2H), 2.36 (s, 3H). MS (m/z): 440 [MH] <sup>+</sup> .
2-2	2-chloro-4-methoxyphenyl	CH <sub>3</sub>	NMR ( <sup>1</sup> H, CDCl <sub>3</sub> ): δ 7.97 (bs, 1H), 7.86 (bs, 1H), 7.36-7.32 (m, 2H), 7.01 (d, 2H), 6.85 (m, 1H), 6.66 (s, 1H), 3.96 (t, 2H), 3.79 (s, 3H), 3.54 (t, 2H), 2.35 (s, 3H). MS (m/z): 424 [MH] <sup>+</sup> .
2-3	2,4-dimethoxyphenyl	CH <sub>3</sub>	NMR ( <sup>1</sup> H, CDCl <sub>3</sub> ): δ 7.97 (d, 1H), 7.86 (d, 1H), 7.34 (d, 1H), 7.04 (d, 1H), 6.64 (s, 1H), 6.54 (d, 1H), 6.51 (d, 1H), 3.82 (s, 3H), 3.79 (s, 3H), 3.53-3.46 (m, 4H), 2.36 (s, 3H). MS (m/z): 421 [MH] <sup>+</sup> .
2-4	2-chloro-4-cyano-phenyl	CH <sub>3</sub>	NMR ( <sup>1</sup> H, CDCl <sub>3</sub> ): δ 8.02 (d, 1H), 7.90 (d, 1H), 7.74 (d, 1H), 7.73 (d, 1H), 7.56 (dd, 1H), 7.37 (d, 1H), 7.09 (d, 1H), 6.83 (s, 1H), 4.16 (t, 2H), 3.62 (t, 2H), 2.43 (s, 3H). MS (m/z): 419 [MH] <sup>+</sup> .

2-5	2-trifluoromethoxy-4-cyanophenyl	CH <sub>3</sub>	NMR ( <sup>1</sup> H, CDCl <sub>3</sub> ): δ 8.66 (d, 1H); 8.03 (s, 1H); 7.96 (d, 1H); 7.94 (d, 1H); 7.88 (dd, 1H); 7.81 (d, 1H); 7.21 (s, 1H); 7.10 (d, 1H); 4.16 (t, 2H); 3.56 (t, 2H); 2.35 (s, 3H). MS (m/z): 469 [MH] <sup>+</sup> .
2-6	2-ethyl-4-cyanophenyl	CH <sub>3</sub>	NMR ( <sup>1</sup> H, DMSO): δ 9.17 (s, 1H), 8.32 (d, 1H), 7.98 (d, 1H), 7.92 (s, 1H), 7.75 (s, 1H), 7.60 (s, 1H), 7.54 (d, 1H), 7.03 (s, 2H), 7.98 (s, 1H), 4.01 (t, 2H), 3.23 (t, 2H), 2.64 (q, 2H), 2.41 (s, 3H), 1.40 (t, 3H). MS (m/z): 414 [MH] <sup>+</sup> .
2-7	2-methyl-4-fluorophenyl	CH <sub>3</sub>	NMR ( <sup>1</sup> H, CDCl <sub>3</sub> ): δ 7.98 (d, 1H), 7.88 (d, 1H), 7.36 (d, 1H), 7.25 (m, 1H), 7.06 (d, 1H), 7.02/6.9 (m, 2H), 6.57 (s, 1H), 3.93 (t, 2H), 3.56 (t, 2H), 2.36 (s, 3H), 2.27 (s, 3H). MS (m/z): 392 [MH] <sup>+</sup> .
2-8	2-methyl-4-(1-pyrazolyl)phenyl	CH <sub>3</sub>	NMR ( <sup>1</sup> H, CDCl <sub>3</sub> ): δ 7.99 (d, 1H), 7.89 (d, 1H), 7.87 (d, 1H), 7.70 (d, 1H), 7.63 (d, 1H), 7.52 (dd, 1H), 7.35 (d, 1H), 7.34 (d, 1H), 7.06 (d, 1H), 6.66 (s, 1H), 6.44 (t, 1H), 3.98 (t, 2H), 3.57 (t, 2H), 2.36 (s, 3H), 2.33 (s, 3H). MS (m/z): 440 [MH] <sup>+</sup> .


## EXAMPLE 3

## Synthesis of representative compounds of structure (VIa)

Example 3-16-Methyl-1-[4-nitro-2-(trifluoromethyl)phenyl]-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine

To a solution of intermediate 32 (17 mg, 0.034 mmol) in anh. CH<sub>2</sub>Cl<sub>2</sub> (5 mL) at r.t., under N<sub>2</sub>, were added Et<sub>3</sub>N (0.019 ml), PPh<sub>3</sub> (17.8 mg) and I<sub>2</sub> (26.1 mg). The reaction mixture was stirred at r.t. for 1 hr. Water (10 mL) was then added and the solution extracted with EtOAc (15 mL). The organic layer was dried over anh. Na<sub>2</sub>SO<sub>4</sub>, the solids were filtered and the solvents evaporated *in vacuo*. The crude compound thus obtained was purified by flash chromatography (silica gel, cHex/EtOAc 6:4) to give 5.9 mg of the title compound as a yellow solid.

All the analytical data are set forth in the following Table 3-1.



Cpd. No.	W1	R <sub>1</sub>	R <sub>4</sub>	Analytical Data
3-1	2-trifluoromethyl-4-nitrophenyl	CH <sub>3</sub>	H	NMR ( <sup>1</sup> H, CDCl <sub>3</sub> ): δ 8.63 (d,1H), 8.41 (dd,1H), 8.03 (d,1H), 7.90 (d,1H), 7.76 (d,1H), 7.37 (d,1H), 7.10 (d,1H), 6.85 (s,1H), 4.08 (t,2H), 3.65 (t,2H), 2.40 (s,3H). MS (m/z): 473 [MH] <sup>+</sup> .

#### EXAMPLE 4

Synthesis of representative compounds of structure (VIIa)

##### Example 4-1

4-[4-(1'-H-1,3'-Bipyrazol-1'-yl)-6-methyl-2,3-dihydro-1H-pyrrolo[2,3-b]pyridin-1-yl]-3-methylbenzonitrile

To a solution of Intermediate 54 (20 mg, 0.049 mmol) in anhyd. CH<sub>2</sub>Cl<sub>2</sub> (2.5 mL), at r.t., under N<sub>2</sub>, were added triphenylphosphine (78 mg, 5 eq) and tetrabromomethane (83 mg 5 eq) and the reaction mixture was stirred at r.t. for 60 min. It was partitioned between CH<sub>2</sub>Cl<sub>2</sub>/sat.aq. NaHCO<sub>3</sub>. The phases were separated and the organic layer was washed with sat.aq. NaCl (2x10 mL). It was dried over anhyd. Na<sub>2</sub>SO<sub>4</sub>, the solids were filtered and the solvent evaporated. The crude product was purified by flash chromatography (silica gel, 7:3 cHex/EtOAc) to give the title compound as a white solid (9 mg, 48%).

##### Example 4-2

4-[4-(1'-H-1,3'-Bipyrazol-1'-yl)-6-methyl-2,3-dihydro-1H-pyrrolo[2,3-b]pyridin-1-yl]-3-(trifluoromethyl)benzonitrile



To a solution of intermediate 55 (85 mg, 0.187 mmol) in anh.  $\text{CH}_2\text{Cl}_2$  (5.5 mL), at r.t., under  $\text{N}_2$ , were added triphenylphosphine (98 mg, 2.5 eq) and tetrabromomethane (124 mg 2.5 eq) and the reaction mixture was stirred at r.t. for 60 min. It was partitioned between  $\text{CH}_2\text{Cl}_2$ /sat.aq.  $\text{NaHCO}_3$ . The phases were separated and the organic layer was washed with sat.aq.  $\text{NaCl}$  (2x10 mL). It was dried over anh.  $\text{Na}_2\text{SO}_4$ , the solids were filtered and the solvent evaporated. The crude product was purified by flash chromatography (silica gel, 7:3 cHex/EtOAc) to give the title compound as a solid (42 mg, 51%).

#### Example 4-3

##### 4-[4-(1'-H-1,3'-Bipyrazol-1'-yl)-6-methyl-2,3-dihydro-1H-pyrrolo[2,3-b]pyridin-1-yl]-3-chloro-benzonitrile


To a solution of intermediate 56 (39 mg, 0.094 mmol) in anh.  $\text{CH}_2\text{Cl}_2$  (5.5 mL), at r.t., under  $\text{N}_2$ , were added triphenylphosphine (49 mg, 2.5 eq) and tetrabromomethane (62 mg 2.5 eq) and the reaction mixture was stirred at r.t. for 60 min. It was partitioned between  $\text{CH}_2\text{Cl}_2$ /sat.aq.  $\text{NaHCO}_3$ . The phases were separated and the organic layer was washed with sat.aq.  $\text{NaCl}$  (2x10 mL). It was dried over anh.  $\text{Na}_2\text{SO}_4$ , the solids were filtered and the solvent evaporated. The crude product was purified by flash chromatography (silica gel, 7:3 cHex/EtOAc) to give the title compound as a solid (14 mg, 35%).

#### Example 4-4

##### 1'-[6-Methyl-1-[2-methyl-4-(methyloxy)phenyl]-2,3-dihydro-1H-pyrrolo[2,3-b]pyridin-4-yl]-1'-H-1,3'-bipyrazole

To a solution of intermediate 84 (200 mg, 0.53 mmol) in anh. NMP (1 mL), at r.t., under  $\text{N}_2$ , were added 3-bromo-1H-pyrazole (77 mg, 1 eq.),  $\text{CuI}$  (101 mg, 1 eq),  $\text{K}_2\text{CO}_3$  (154 mg, 2.1 eq) and (1R,2R)-diaminomethylcyclohexane (23 mg, 0.3 eq.). The reaction mixture was heated at  $150^\circ\text{C}$  for 3.5 h. It was cooled down to r.t., poured into water/EtOAc and the phases were separated. The aqueous layer was further extracted with EtOAc (2x10 mL). The combined organic extracts were dried over anh.  $\text{Na}_2\text{SO}_4$ , the solids were filtered and the solvent evaporated. The residue was purified by flash chromatography (silica gel, 95:5 cHex/EtOAc, two purifications) to give the title compound as a white solid (7.5 mg, 4%).

All the analytical data are set forth in the following Table 4-1.



Cpd. No.	W1	R <sub>1</sub>	Analytical Data
4-1	2-methyl-4-cyano-phenyl	CH <sub>3</sub>	NMR ( <sup>1</sup> H, CDCl <sub>3</sub> ): δ 8.2 (d, 1H), 8.0 (d, 1H), 7.7 (d, 1H), 7.6 (d, 1H), 7.5 (dd, 1H), 7.4 (dd, 1H), 6.8 (d, 1H), 6.7 (d, 1H), 6.4 (t, 1H), 4.0 (t, 2H), 4.0 (t, 2H), 3.6 (t, 2H), 2.41, 2.35 (s+s, 6H). MS (m/z): 382 [MH] <sup>+</sup> .
4-2	2-trifluoromethyl-4-cyanophenyl	CH <sub>3</sub>	NMR ( <sup>1</sup> H, CDCl <sub>3</sub> ): δ 8.2 (d, 1H), 8.0 (d+d, 2H), 7.8 (dd, 1H), 7.7 (d, 1H), 7.6 (dd, 1H), 6.8 (m, 2H), 6.4 (d, 1H), 4.0 (t, 2H), 3.6 (t, 2H), 2.41 (s, 3H). MS (m/z): 436 [MH] <sup>+</sup> .
4-3	2-chloro-4-cyano-phenyl	CH <sub>3</sub>	NMR ( <sup>1</sup> H, CDCl <sub>3</sub> ): δ 8.2 (d, 1H), 8.0 (d, 1H), 7.8 (m, 3H), 7.6 (d, 1H), 6.8 (m, 2H), 6.4 (d, 1H), 4.2 (t, 2H), 3.6 (t, 2H), 2.40 (s, 3H). MS (m/z): 402 [MH] <sup>+</sup> .
4-4	2-methyl-4-methoxy-phenyl	CH <sub>3</sub>	NMR ( <sup>1</sup> H, Acetone-d <sub>6</sub> ): δ 8.15 (d, 1H), 7.86 (d, 1H), 7.66 (d, 1H), 7.12 (d, 1H), 6.76 (d, 1H), 6.72 (dd, 1H), 6.54 (s, 1H), 6.39 (dd, 1H), 3.84 (t, 2H), 3.74 (s, 3H), 3.44 (t, 2H), 2.28 (s, 3H), 2.18 (s, 3H). MS (m/z): 387 [MH] <sup>+</sup> .

## EXAMPLE 5

## Synthesis of representative compounds of structure (VIIIa)

Example 5-13-Methyl-4-[6-methyl-4-[3-(2-pyridinyl)-1H-pyrazol-1-yl]-2,3-dihydro-1H-pyrrolo[2,3-b]pyridin-1-yl]benzonitrile

To a solution of intermediate 69 (24 mg, 0.058 mmol) in anh. CH<sub>2</sub>Cl<sub>2</sub> (2 mL), at r.t., under N<sub>2</sub>, were added PPh<sub>3</sub> (30 mg, 2 eq) and CBr<sub>4</sub> (29 mg, 1.5 eq) and the reaction mixture was stirred at r.t. for 18 hr. It was partitioned between CH<sub>2</sub>Cl<sub>2</sub>/sat.aq. NaHCO<sub>3</sub>. The phases were separated and the organic layer was washed with sat.aq. NaCl (2x10 mL). It was dried over anh. Na<sub>2</sub>SO<sub>4</sub>, the solids were filtered and the solvent evaporated. The crude product was purified by flash chromatography (silica gel, cHex/EtOAc 7:3) to give the title compound as a solid (8 mg, 35%).


Example 5-23-Chloro-4-[6-methyl-4-[3-(2-pyridinyl)-1H-pyrazol-1-yl]-2,3-dihydro-1H-pyrrolo[2,3-b]pyridin-1-yl]benzonitrile

To a solution of intermediate 70 (39 mg, 0.094 mmol) in anh. CH<sub>2</sub>Cl<sub>2</sub> (5 mL), at r.t., under N<sub>2</sub>, were added imidazole (18 mg, 3 eq), triphenylphosphine (47 mg, 2 eq) and iodine (69.3 mg, 3 eq) and the reaction mixture was stirred at r.t. for 1 hr. It was then partitioned between EtOAc/sat.aq. NaCl. The phases were separated and the organic layer was washed with sat.aq. NaCl (2x10 mL). It was dried over anh. Na<sub>2</sub>SO<sub>4</sub>, the solids were filtered and the solvent evaporated. The crude product was purified by flash chromatography (silica gel, 1:1 cHex/EtOAc) to give the title compound as a colorless oil (4.5 mg, 12%).

Example 5-34-[6-Methyl-4-[3-(2-pyridinyl)-1H-pyrazol-1-yl]-2,3-dihydro-1H-pyrrolo[2,3-b]pyridin-1-yl]-3-(trifluoromethyl)benzonitrile

To a solution of intermediate 71 (44 mg, 0.095 mmol) in anh. CH<sub>2</sub>Cl<sub>2</sub> (3 mL), at r.t., under N<sub>2</sub>, were added PPh<sub>3</sub> (50 mg, 2 eq) and CBr<sub>4</sub> (47 mg, 1.5 eq) and the reaction mixture was stirred at r.t. for 18 hr. It was partitioned between CH<sub>2</sub>Cl<sub>2</sub>/sat.aq. NaHCO<sub>3</sub>. The phases were separated and the organic layer was washed with sat.aq. NaCl (2x10 mL). It was dried over anh. Na<sub>2</sub>SO<sub>4</sub>, the solids were filtered and the solvent evaporated. The crude product was purified by flash chromatography (silica gel, cHex/EtOAc 1:1) to give the title compound as a solid (8 mg, 20%).

All the analytical data are set forth in the following Table 5-1.



Cpd. No.	W1	R1	Analytical Data
5-1	2-methyl-4-cyano-phenyl	CH <sub>3</sub>	NMR ( <sup>1</sup> H, CDCl <sub>3</sub> ): δ 8.65 (dd, 1H), 8.10 (d, 1H), 8.00 (d, 1H), 7.75 (dt, 1H), 7.55-7.48 (m, 2H), 7.39 (d, 1H), 7.26 (m, 1H), 7.15 (d, 1H), 6.78 (s, 1H), 4.00 (t, 2H), 3.60 (t, 2H), 2.37 (s, 3H), 2.31 (s, 3H). MS (m/z): 393 [MH] <sup>+</sup> .
5-2	2-chloro-4-cyano-phenyl	CH <sub>3</sub>	NMR ( <sup>1</sup> H, CDCl <sub>3</sub> ): δ 8.67 (d, 1H), 8.11 (d, 1H), 8.03 (d, 1H), 7.75 (m, 2H), 7.75 (m, 1H), 7.55 (dd, 1H), 7.26 (m, 1H), 7.2 (d, 1H), 6.88 (s, 1H), 4.16 (t, 2H), 3.65 (t, 2H), 2.43 (s, 3H). MS (m/z): 413 [MH] <sup>+</sup> .
5-3	2-trifluoromethyl-4-cyano-phenyl	CH <sub>3</sub>	NMR ( <sup>1</sup> H, CDCl <sub>3</sub> ): δ 8.66 (dd, 1H), 8.10 (d, 1H), 8.02 (d, 1H), 8.00 (d, 1H), 7.84 (dd, 1H), 7.76 (t, 1H), 7.69 (d, 1H), 7.27 (t, 1H), 7.18 (d, 1H), 6.87 (s, 1H), 4.02 (t, 2H), 3.64 (t, 2H), 2.38 (s, 3H). MS (m/z): 447 [MH] <sup>+</sup> .

#### EXAMPLE 6

##### CRF Binding Activity

CRF binding affinity has been determined in vitro by the compounds' ability to displace <sup>125</sup>I-oCRF and <sup>125</sup>I-Sauvagine for CRF1 and CRF2 SPA, respectively, from recombinant human CRF receptors expressed in Chinese Hamster Ovary (CHO) cell membranes. For membrane preparation, CHO cells from confluent T-flasks were collected in SPA buffer (HEPES/KOH 50mM, EDTA 2mM; MgCl<sub>2</sub> 10mM, pH 7.4.) in 50mL centrifuge tubes, homogenized with a

Polytron and centrifuged (50'000g for 5min at 4°C: Beckman centrifuge with JA20 rotor). The pellet was resuspended, homogenized and centrifuged as before.

The SPA experiment has been carried out in Optiplate by the addition of 100 µL the reagent mixture to 1µL of compound dilution (100% DMSO solution) per well. The assay mixture was prepared by mixing SPA buffer, WGA SPA beads (2.5 mg/mL), BSA (1 mg/mL) and membranes (50 and 5 µg of protein/mL for CRF1 and CRF2 respectively) and 50 pM of radioligand.

The plate was incubated overnight (>18 hrs) at room temperature and read with the Packard Topcount with a WGA-SPA <sup>125</sup>I counting protocol.

#### EXAMPLE 7

##### CRF functional assay

Compounds of the invention were characterised in a functional assay for the determination of their inhibitory effect. Human CRF-CHO cells were stimulated with CRF and the receptor activation was evaluated by measuring the accumulation of cAMP.

CHO cells from a confluent T-flask were resuspended with culture medium without G418 and dispensed in a 96-well plate, 25'000c/well, 100 µL/well and incubated overnight. After the incubation the medium was replaced with 100 µL of cAMP IBMX buffer warmed at 37°C (5mM KCl, 5mM NaHCO<sub>3</sub>, 154mM NaCl, 5mM HEPES, 2.3mM CaCl<sub>2</sub>, 1mM MgCl<sub>2</sub>, 1g/L glucose, pH 7.4 additioned by 1mg/mL BSA and 1mM IBMX) and 1µL of antagonist dilution in neat DMSO. After 10 additional minutes of incubation at 37°C in a plate incubator without CO<sub>2</sub>, 1µL of agonist dilution in neat DMSO was added. As before, the plate was incubated for 10 minutes and then cAMP cellular content was measured by using the Amersham RPA 538 kit.


All publications, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference as if each individual publication were specifically and individually indicated to be incorporated by reference herein as though fully set forth.

It is to be understood that the present invention covers all combinations of particular and preferred groups described herein above.

The application of which this description and claims forms part may be used as a basis for priority in respect of any subsequent application. The claims of such subsequent application may be directed to any feature or combination of features described herein. They may take the form of product, composition, process, or use claims and may include, by way of example and without limitation, the following claims:

## Claims

1. Compounds of formula (I) including stereoisomers, prodrugs and pharmaceutically acceptable salts or solvates thereof




wherein

- R is aryl or heteroaryl, each of which may be substituted by 1 or more Z groups;
- R<sub>1</sub> is hydrogen, C3-C7 cycloalkyl, C1-C6 alkyl, C1-C6 alkoxy, C1-C6 thioalkyl, C2-C6 alkenyl, C2-C6 alkynyl, halo C1-C6 alkyl, halo C1-C6 alkoxy, halogen, NR<sub>6</sub>R<sub>7</sub> or cyano;
- R<sub>2</sub> and R<sub>3</sub> together with N form a 5-6 membered aromatic heterocycle, which is substituted by at least one group R<sub>8</sub> and may be further substituted by 1 to 3 R<sub>8</sub> groups;
- R<sub>4</sub> is hydrogen, C1-C6 alkyl, halogen or halo C1-C6 alkyl;
- R<sub>5</sub> is a C1-C4 alkyl, -OR<sub>6</sub> or -NR<sub>6</sub>R<sub>7</sub>;
- R<sub>6</sub> is hydrogen or C1-C6 alkyl;
- R<sub>7</sub> is hydrogen or C1-C6 alkyl;
- R<sub>8</sub> is a 5-6 membered aromatic heterocycle, which may be substituted by 1 to 4 R<sub>10</sub> groups;
- R<sub>9</sub> is hydrogen, C3-C7 cycloalkyl, C3-C7 cycloalkenyl, C1-C6 alkyl, C1-C6 alkoxy, halo C1-C6 alkyl, halo C1-C6 alkoxy, hydroxy, halogen, nitro, cyano, -C(O)R<sub>5</sub>, C(O)NR<sub>6</sub>R<sub>7</sub>, phenyl which may be substituted by 1 to 4 R<sub>10</sub> groups;
- R<sub>10</sub> is C1-C6 alkyl, halo C1-C2 alkyl, halogen, nitro, hydroxy, halo C1-C6 alkoxy, C1-C6 alkoxy, or cyano;
- Z is selected in a group consisting from halogen, C1-C6 alkyl, C1-C6 alkoxy, halo C1-C6 alkyl, C2-C6 alkenyl, C2-C6 alkynyl, halo C1-C6 alkoxy, -C(O)R<sub>5</sub>, -NR<sub>6</sub>R<sub>7</sub>, nitro, cyano, and a group R<sub>8</sub>;
- and when R<sub>4</sub> is hydrogen, R<sub>2</sub> and R<sub>3</sub> together with N form a pyrazolyl group substituted with at least one R<sub>8</sub> corresponding to a thiazolyl group, and R corresponds to a phenyl group, and the substituent Z is at least one nitro group,
- then in the compounds of formula (I) the nitro group is not present in the ortho position with respect to the nitrogen atom present in the 5-membered ring, named as B;


and the compounds of formula (I) don't include the following:

- 1-(2,4-bis-trifluoromethyl-phenyl)-6-methyl-4-(3-thiazol-2-yl-pyrazol-1-yl)-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine;
- 3-methyl-4-[6-methyl-4-(3-thiazol-2-yl-pyrazol-1-yl)-2,3-dihydro-1H-pyrrolo[2,3-b]pyridin-1-yl]-benzonitrile;
- 4-[6-methyl-4-(3-thiazol-2-yl-pyrazol-1-yl)-2,3-dihydro-1H-pyrrolo[2,3-b]pyridin-1-yl]-3-trifluoromethyl-benzonitrile;
- 6-methyl-1-(2-methyl-4-trifluoromethoxy-phenyl)-4-(3-thiazol-2-yl-pyrazol-1-yl)-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine;
- 1-(4-methoxy-2-methyl-phenyl)-6-methyl-4-(3-thiazol-2-yl-pyrazol-1-yl)-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine
- 1-(2,4-bis-trifluoromethyl-phenyl)-6-methyl-4-(3-pyridin-2-yl-pyrazol-1-yl)-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine
- 4-[1,3'bipyrazolyl-1'-yl-1-(2,4-bis-trifluoromethyl-phenyl)-6-methyl-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine.

2. Compounds of formula (II), according to claim 1, in which  $R_4$  is hydrogen and  $R$ ,  $R_1$ ,  $R_2$ , and  $R_3$  are defined as in claim 1.




3. Compounds of formula (III), according to claim 2,



in which

$n$  is an integer from 1 to 2; and  
 $R$ ,  $R_1$ , and  $R_2$  and  $R_3$  are defined as in claim 1.

4. Compounds of formula (IV), according to claim 3,



In which:

$R_9$  is a thiazolyl derivative;


$n$  is an integer from 1 to 2;

$R$  corresponds to  $W$  and

$W$  is a pyridine derivative, which may be substituted by 1 to 4  $Z$  groups, as defined above; and


$R_1$ ,  $R_9$  and  $R_{10}$  are defined as in claim 1.

5. Compounds of formula (IVa), according to claim 4,



in which  $W$  is defined as in claim 4.

6. Compounds of formula (V), according to claim 3,





in which:

$R_8$  is a thiazolyl derivative,;

$n$  is an integer from 1 to 2;


$R$  corresponds to  $W_1$  and

$W_1$  is a phenyl derivative substituted by 2 to 4  $Z_1$  groups;

$Z_1$  is selected in a group consisting from halogen, C1-C6 alkyl, C1-C6 alkoxy, halo C1-C6 alkyl, C2-C6 alkenyl, C2-C6 alkynyl, halo C1-C6 alkoxy,  $-C(O)R_8$ ,  $-NR_8R_7$ , cyano, and a group  $R_8$ ;


and  $R_1$ ,  $R_5$ ,  $R_6$ ,  $R_7$ ,  $R_8$ ,  $R_9$  and  $R_{10}$  are defined as in claim 1.

7. Compounds of formula (Va), according to claim 6,



in which  $W_1$  is defined as in claim 6.

8. Compounds of formula (VI), according to claim 3,



in which:

$R_8$  is a thiazolyl derivative;


$n$  is an integer from 1 to 2;

$R$  corresponds to  $W_2$  and

$W_2$  is a phenyl derivative, which may be substituted by 2 to 4  $Z$  groups as defined in claim 1 provided that at least one  $Z$  group is nitro and further provided that the nitro group is not in the ortho position with respect to the nitrogen atom of the 5-membered ring named as B; and


$R_1$ ,  $R_9$  and  $R_{10}$  are defined as in claim 1.

9. Compounds of formula (VIa), according to claim 8,



In which  $W_2$  is defined as in claim 8.

10. Compounds of formula (VII), according to claim 3,




in which

$R_8$  is a pyrazolyl derivative;

$n$  is an integer from 1 to 2;


$R$ ,  $R_9$  and  $R_{10}$  are defined as in claim 1.

11. Compounds of formula (VIIa), according to claim 10,



in which  $W_1$  is defined as in claim 6.


12. Compounds of formula (VIII), according to claim 3,



In which

- $R_9$  is a pyridine derivative;  
 $n$  is an integer from 1 to 2;  
 $m$  is an integer from 1 to 3;  
 $R$ ,  $R_9$  and  $R_{10}$  are defined as in claim 1.

13. Compounds of formula (VIIIa), according to claim 12,




in which  $W_1$  is defined as in claim 6.

14. Compounds according to the claims from 1 to 13 selected in the group consisting from:  
 6-methyl-1-[6-(methyloxy)-2-(trifluoromethyl)-3-pyridinyl]-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine;  
 6-methyl-1-[2-methyl-6-(methyloxy)-3-pyridinyl]-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine;  
 1-[2,6-bis(methyloxy)-3-pyridinyl]-6-methyl-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine;  
 N,N,4-trimethyl-5-[6-methyl-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-2,3-dihydro-1H-pyrrolo[2,3-b]pyridin-1-yl]-2-pyridinamine;  
 1-(2-difluoromethyl-4-methoxy-phenyl)-6-methyl-4-(3-thiazol-2-yl-pyrazol-1-yl)-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine;

1-(2-chloro-4-methoxy-phenyl)-6-methyl-4-(3-thiazol-2-yl-pyrazol-1-yl)-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine;  
1-[2,4-bis(methyloxy)phenyl]-6-methyl-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine;  
3-chloro-4-{6-methyl-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-2,3-dihydro-1H-pyrrolo[2,3-b]pyridin-1-yl}benzonitrile;  
4-{6-methyl-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-2,3-dihydro-1H-pyrrolo[2,3-b]pyridin-1-yl}-3-[(trifluoromethyl)oxy]benzonitrile;  
3-ethyl-4-{6-methyl-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-2,3-dihydro-1H-pyrrolo[2,3-b]pyridin-1-yl}benzonitrile;  
1-(4-fluoro-2-methylphenyl)-6-methyl-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine;  
6-methyl-1-[2-methyl-4-(1H-pyrazol-1-yl)phenyl]-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine;  
6-methyl-1-[4-nitro-2-(trifluoromethyl)phenyl]-4-[3-(1,3-thiazol-2-yl)-1H-pyrazol-1-yl]-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine;  
4-[4-(1H-1,3'-bipyrazol-1'-yl)-6-methyl-2,3-dihydro-1H-pyrrolo[2,3-b]pyridin-1-yl]-3-methylbenzonitrile;  
4-[4-(1H-1,3'-bipyrazol-1'-yl)-6-methyl-2,3-dihydro-1H-pyrrolo[2,3-b]pyridin-1-yl]-3-(trifluoromethyl)benzonitrile;  
4-[4-(1H-1,3'-bipyrazol-1'-yl)-6-methyl-2,3-dihydro-1H-pyrrolo[2,3-b]pyridin-1-yl]-3-chlorobenzonitrile;  
1'-{6-methyl-1-[2-methyl-4-(methyloxy)phenyl]-2,3-dihydro-1H-pyrrolo[2,3-b]pyridin-4-yl}-1H-1,3'-bipyrazole;  
3-methyl-4-{6-methyl-4-[3-(2-pyridinyl)-1H-pyrazol-1-yl]-2,3-dihydro-1H-pyrrolo[2,3-b]pyridin-1-yl}benzonitrile;  
3-chloro-4-{6-methyl-4-[3-(2-pyridinyl)-1H-pyrazol-1-yl]-2,3-dihydro-1H-pyrrolo[2,3-b]pyridin-1-yl}benzonitrile;  
4-{6-methyl-4-[3-(2-pyridinyl)-1H-pyrazol-1-yl]-2,3-dihydro-1H-pyrrolo[2,3-b]pyridin-1-yl}-3-(trifluoromethyl)benzonitrile.

15. A process for the preparation of a compound of formula (I) as claimed in claim 1, which comprises the following steps:



in which

- step a stands for conversion of the leaving group L, selected in a group consisting from: halogen or reactive residue of sulphonic acid (e.g. mesylate, tosylate), preferably chloride, in the amino group of compounds (IV), by reaction with the suitable amine  $\text{NR}_2\text{R}_3$  in basic conditions;
- step b stands for reduction of the ester group with a suitable reducing agent (such as DIBAL-H) to hydroxy group of compounds (V);
- step c stands for oxidation of the hydroxy group with a suitable oxidising agent (such as Dess-Martin periodinane) to aldehyde group of compound (VI);
- step d stands for formation of the aldehyde group of compounds (VIII) by Wittig reaction in the usual conditions, through formation of enol ether followed by acid hydrolysis (step e);
- step f stands for reduction of the aldehyde group with a suitable reducing agent (such as  $\text{NaBH}_4$ ) to hydroxy group of compounds (IX);

- step g stands for conversion of the hydroxy group in the suitable protecting group of compounds (X) (such as TBS: *tert*-butyldimethylsilyl);
- step h stands for Buchwald reaction by coupling with the suitable amine RNH<sub>2</sub>;
- step i stands for deprotection reaction to give the hydroxy group of compounds (XII);
- step l stands for intramolecular cyclisation by heating after conversion of the hydroxy group of compounds (XII) in a suitable leaving group (such as bromide, by reaction with CBr<sub>4</sub> and PPh<sub>3</sub>) to give the final compounds (IIa).
16. The use of a compound according to anyone of claims from 1 to 14, in the preparation of a medicament for use in the treatment of conditions mediated by CRF (corticotropin-releasing factor).
17. The use of a compound according to anyone of claims from 1 to 14, in the preparation of a medicament for use in the treatment of depression and anxiety.
18. The use of a compound according to anyone of claims from 1 to 14, in the preparation of a medicament for use in the treatment of IBS (irritable bowel disease) and IBD (inflammatory bowel disease).
19. A compound according to anyone of claims from 1 to 14, for use in the treatment of conditions mediated by CRF (corticotropin-releasing factor).
20. A pharmaceutical composition comprising a compound of claims from 1 to 14, in admixture with one or more physiologically acceptable carriers or excipients.
21. A method for the treatment of a mammal, including man, in particular in the treatment of conditions mediated by CRF (corticotropin-releasing factor), comprising administration of an effective amount of a compound of anyone of claims from 1 to 14.

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP2004/000409

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/437 A61P1/00 A61P25/18 A61P25/22 A61P25/24  
A61P29/00 C07D471/04

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K A61P C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BEILSTEIN Data, BIOSIS, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	WO 03/008412 A (GLAXO GROUP LTD ; ST-DENIS YVES (IT); DI FABIO ROMANO (IT); MICHELI FA) 30 January 2003 (2003-01-30) cited in the application Claims, examples	1-21
A	WO 95/33750 A (CHEN YUHPYNG L ; PFIZER (US)) 14 December 1995 (1995-12-14) cited in the application the whole document	1-21
A	WO 98/45295 A (HORVATH RAYMOND F ; NEUROGEN CORP (US); HUTCHINSON ALAN (US)) 15 October 1998 (1998-10-15) the whole document	1-21

☐ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document but published on or after the international filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*8\* document member of the same patent family

Date of the actual completion of the international search

26 May 2004

Date of mailing of the international search report

04/06/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel: (+31-70) 340-2040, Tx: 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Stroeter, T

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP2004/000409

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 03008412	A	30-01-2003	CA 2451530 A1 WO 03008412 A2 GB 2378702 A	30-01-2003 30-01-2003 19-02-2003
WO 9533750	A	14-12-1995	AT 196295 T AU 692548 B2 AU 2453095 A BR 9502708 A CA 2192354 A1 CN 1150428 A ,B CN 1246475 A CZ 9603608 A3 DE 69518841 D1 DE 69518841 T2 DK 764166 T3 EP 0764166 A1 ES 2150567 T3 FI 964894 A GR 3034765 T3 HR 950321 A1 HU 75774 A2 WO 9533750 A1 JP 2000001434 A JP 3223169 B2 JP 11246411 A JP 9507249 T JP 3193055 B2 NO 965237 A NO 2391 A NZ 285442 A PL 320631 A1 PT 764166 T RU 2221799 C2 SK 155596 A3 TW 530047 B US 5962479 A ZA 9504677 A	15-09-2000 11-06-1998 04-01-1996 30-04-1996 14-12-1995 21-05-1997 08-03-2000 14-07-1999 19-10-2000 11-01-2001 09-10-2000 26-03-1997 01-12-2000 05-12-1996 28-02-2001 28-02-1998 28-05-1997 14-12-1995 07-01-2000 29-10-2001 14-09-1999 22-07-1997 30-07-2001 06-02-1997 06-02-1997 27-05-1998 13-10-1997 31-01-2001 20-01-2004 11-12-2000 01-05-2003 05-10-1999 09-12-1996
WO 9845295	A	15-10-1998	CA 2207784 A1 US 5955613 A AU 2622797 A WO 9845295 A1 AU 738304 B2 US 2002111490 A1 US 6133282 A US 6355651 B1	13-12-1998 21-09-1999 29-10-1998 15-10-1998 13-09-2001 15-08-2002 17-10-2000 12-03-2002